

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

09/857372

INTERNATIONAL APPLICATION NO.

PCT/EP99/09833

INTERNATIONAL FILING DATE

December 6, 1999

PRIORITY DATE CLAIMED

December 4, 1998

TITLE OF INVENTION

DRUG TARGETS IN CANDIDA ALBICANS


APPLICANT(S) FOR DO/EO/US :

CONTRERAS

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.
 2. ☐ This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371.
 3. ☐ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
 4. ☐ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
 5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☒ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
 6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
 7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☐ have not been made and will not be made.
 8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
 9. ☒ An executed oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
 10. ☐ A translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).
- Items 11. to 16. below concern document(s) or information included:
11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
 12. ☒ Assignment documents (3) for recording. Separate cover sheets in compliance with 37 CFR 3.28 and 3.31 are included.
 13. ☐ A FIRST preliminary amendment.
☐ A SECOND or SUBSEQUENT preliminary amendment.
 14. ☒ A substitute specification in 71 pages including 30 claims, a sequence listing, and 4 figures.
 15. ☐ A change of power of attorney and/or address letter.
 16. ☒ Other items or information: Copy of the International Preliminary Examination Report; Copy of the International Search Report; Associate Power of Attorney; Sequence Disk; Sequence Listing, Verified Statement.

JC12 Rec-0 CT/PTO 04 JUN 2001

U.S. APPLICATION NO. (if known, see 37 CFR 1.5)		INTERNATIONAL APPLICATION NO.		ATTORNEY'S DOCKET NUMBER
09/857372		PCT/EP99/09833		JAB-1430
17. <input checked="" type="checkbox"/> The following fees are submitted:				CALCULATIONS PTO USE ONLY
BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)):				
Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO..... \$1070.00				
International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO.....\$930.00				
International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.455(a)(2)) paid to USPTO..... \$790.00				
International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4)..... \$720.00				
International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4)..... \$98.00				
ENTER APPROPRIATE BASIC FEE AMOUNT =				\$ 930.00
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	
Total claims	30 - 20 =	10	x \$18.00	\$180.00
Independent claims	20 - 3 =	17	x \$80.00	\$1360.00
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$270.00	\$
TOTAL OF ABOVE CALCULATIONS =				\$2470.00
Reduction of 1/2 for filing by small entity, if applicable. A Small Entity Statement must also be filed (Note 37 CFR 1.9, 1.27, 1.28).				\$
SUBTOTAL =				\$
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				\$
TOTAL NATIONAL FEE =				\$2470.00
Fee for recording the enclosed 3 assignments (37 CFR 1.21(h)). The assignments must be accompanied by an appropriate cover sheets (37 CFR 3.28, 3.31). \$40.00 per property +				\$120.00
TOTAL FEES ENCLOSED =				\$2470.00
				Amount to be refunded:
				\$
				charged:
				\$ 2590.00
a. <input type="checkbox"/> A check in the amount of \$_____ to cover the above fees is enclosed.				
b. <input checked="" type="checkbox"/> Please charge my Deposit Account No. <u>10-0750/JAB-1430/MHM</u> in the amount of \$2270.00 to cover the above fees. A duplicate copy of this sheet is enclosed.				
c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>10-0750/JAB-1430/MHM</u> . A duplicate copy of this sheet is enclosed.				
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137 (a) or (b)) must be filed and granted to restore the application to pending status.				
SEND ALL CORRESPONDENCE TO:				
Philip S. Johnson, Esq. Johnson & Johnson One Johnson & Johnson Plaza New Brunswick, NJ 08933-7003 USA				
 Signature Myra H. McCormack Reg. No. 36,602 Attorney for Applicants June 4, 2001				

OPPTS - 1 -
DRUG TARGETS IN CANDIDA ALBICANS

Field of the Invention

The present invention is concerned with the
5 identification of genes or functional fragments
thereof from *Candida albicans* which are critical for
growth and cell division and which genes may be used
as selective drug targets to treat *Candida albicans*
associated infections. Novel nucleic acid sequences
10 from *Candida albicans* are also provided and which
encode the polypeptides, which are critical for growth
of *Candida albicans*.

Background of the Invention

15 Opportunistic infections in immunocompromised
hosts represent an increasingly common cause of
mortality and morbidity. *Candida* species are among
the most commonly identified fungal pathogens
associated with such opportunistic infections, with
20 *Candida albicans* being the most common species. Such
fungal infections are thus problematical in, for
example, AIDS populations in addition to normal
healthy women where *Candida albicans* yeasts represent
the most common cause of vulvovaginitis.

25 Although compounds do exist for treating such
disorders, such as, amphotericin, these drugs are
generally limited in their treatment because of their
toxicity and side effects. Therefore, there exists a
need for new compounds, which may be used to treat
30 *Candida* associated infections in addition to
compounds, which are selective in their action against
Candida albicans.

Classical approaches for identifying anti-fungal
compounds have relied almost exclusively on inhibition
35 of fungal or yeast growth as an endpoint. Libraries
of natural products, semi-synthetic, or synthetic

chemicals are screened for their ability to kill or arrest growth of the target pathogen or a related nonpathogenic model organism. These tests are cumbersome and provide no information about a
5 compounds mechanism of action. The promising lead compounds that emerge from such screens must then be tested for possible host-toxicity and detailed mechanism of action studies must subsequently be conducted to identify the affected molecular target.

10

Summary of the Invention

The present inventors have now identified a range of nucleic acid sequences from *Candida albicans*, which
15 encode polypeptides, which are critical for its survival and growth. These sequences represent novel targets which can be incorporated into an assay to selectively identify compounds capable of inhibiting expression of such polypeptides and their potential
20 use in alleviating diseases or conditions associated with *Candida albicans* infection.

In a first aspect of the invention, the invention relates to a nucleic acid molecule encoding a polypeptide which is critical for survival and growth
25 of the yeast *Candida albicans* and which nucleic acid molecule is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 9. More preferably where nucleic acid molecule is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 3 and still
30 more preferably consisting of SEQ ID NO:1 and SEQ ID NO:2.

In a preferred embodiment of this aspect of the invention, the nucleic acid molecule is either RNA or DNA, and in one embodiment, cDNA. The invention also
35 relates to a first nucleic acid molecule capable of hybridising to a second nucleic acid molecule encoding a polypeptide which is critical for survival and

growth of the yeast *Candida albicans* and which second nucleic acid molecule is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 9 under high stringency conditions. In yet another embodiment, the invention relates to an antisense molecule comprising a first nucleic acid molecule capable of hybridising to a second nucleic acid molecule encoding a polypeptide which is critical for survival and growth of the yeast *Candida albicans* wherein the second nucleic acid molecule is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 9.

The invention also relates to cells containing a nucleic acid molecule encoding a polypeptide which is critical for survival and growth of the yeast *Candida albicans* and which nucleic acid molecule is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 9, wherein said cells are bacterial or eukaryotic. The invention further relates to polypeptide encoded by a nucleic acid molecule which is critical for survival and growth of the yeast *Candida albicans* and which nucleic acid molecule is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 9 and to polypeptide having any of amino acid sequences selected from the group consisting of SEQ ID NO: 10 to SEQ ID NO: 16.

The invention also relates to a recombinant DNA construct comprising a nucleic acid molecule encoding a polypeptide which is critical for survival and growth of the yeast *Candida albicans* and which nucleic acid molecule is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 9, wherein the nucleic acid molecule is DNA. Preferably the DNA is an expression vector and preferably the expression vector includes an inducible promoter and/or a reporter molecule. The invention also relates to recombinant DNA constructs comprising a first nucleic acid molecule wherein the first nucleic acid molecule or a molecule

complementary to the first nucleic acid molecule is capable of hybridising to a second nucleic acid molecule encoding a polypeptide which is critical for survival and growth of the yeast *Candida albicans* and
5 which second nucleic acid molecule is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 9 wherein the first nucleic acid molecule is inserted in the antisense orientation.

The invention also relates to cells containing a
10 recombinant DNA construct comprising a nucleic acid molecule encoding a polypeptide which is critical for survival and growth of the yeast *Candida albicans* and which nucleic acid molecule is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 9, wherein
15 said cells are bacterial or eukaryotic.

In another aspect of this invention, the nucleic acid molecules or polypeptides of this invention are provided in a pharmaceutically acceptable carrier, diluent or excipient.

20 The invention further relates to a method for treating a *Candida albicans*-associated disease comprising the step of: administering a composition of matter comprising an antisense nucleic acid molecule capable of binding to a *Candida albicans*-originating
25 nucleic acid sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:9.

In another aspect of this invention, the invention relates to a *Candida albicans* cell comprising an induced mutation in the DNA sequence
30 encoding a polypeptide which is critical for survival and growth of the yeast *Candida albicans* wherein the polypeptide is encoded by a nucleic acid molecule is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 9.

35 The invention also relates to a method of identifying compounds which selectively modulate expression or functionality of polypeptides or

metabolic pathways in which these polypeptides are involved and which polypeptides are crucial for growth and survival of *Candida albicans*, which method comprises: contacting a compound to be tested with one
5 or more *Candida albicans* cells having a mutation in a nucleic acid molecule, the nucleic acid molecule encoding a polypeptide which is critical for survival and growth of the yeast *Candida albicans* and which nucleic acid molecule is selected from the group
10 consisting of SEQ ID NO: 1 to SEQ ID NO: 9 wherein the mutation results in overexpression or underexpression of said polypeptide in addition to contacting one or more wild type *Candida albicans* cells with said compound; and monitoring the growth and/or activity of
15 said mutated cell compared to said wild type; wherein differential growth or activity of said one or more mutated *Candida* cells is indicative of selective action of said compound on a polypeptide or another polypeptide in the same or a parallel pathway. The
20 invention further relates to compounds identified by this method. The compounds can be provided in a pharmaceutical composition with a pharmaceutically acceptable carrier, diluent or excipient therefor.

The invention also relates to a method of
25 identifying DNA sequences from a cell or organism which DNA encodes polypeptides which are critical for growth or survival of said cell or organism, which method comprises: preparing a cDNA or genomic library from said cell or organism in a suitable expression
30 vector which vector is such that it can either integrate into the genome in said cell or that it permits transcription of antisense RNA from the nucleotide sequences in said cDNA or genomic library; selecting transformants exhibiting impaired growth and
35 determining the nucleotide sequence of the cDNA or genomic sequence from the library included in the vector from said transformant. In a preferred aspect

of this invention, the cell or organism is a yeast or filamentous fungus and more preferably the cell or organism is any of *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe* or *Candida albicans*.

5 The invention further relates to antibody capable of specifically binding to a polypeptide of this invention. The invention also relates to oligonucleotides comprising a fragment of from 10 to 120 contiguous nucleotides of a nucleic acid molecule
10 encoding a polypeptide which is critical for survival and growth of the yeast *Candida albicans* and which nucleic acid molecule is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 9. Preferably the oligonucleotide comprises a fragment of
15 from 10 to 50 contiguous nucleotides.

Brief Description of the Figures

Figure 1 is a diagrammatic representation of plasmid pGAL1PNiST-1.

20 Figure 2 is a diagrammatic representation of plasmid pGAL1PSiST-1.

Figure 3 is an exemplary growth curve of a *Candida albicans* strain demonstrating antisense-induced reduction in growth in response to
25 intracellular antisense expressing specifically targeting one of the nucleic acid sequences of this invention.

Figure 4 is a representative growth curve of a *Candida albicans* knock out strain as compared to wild
30 type *Candida albicans*.

Detailed Description of the Invention

According to a first aspect of the invention there is provided a nucleic acid molecule encoding a
35 polypeptide which is critical for survival and growth of the yeast *Candida albicans* and which nucleic acid

molecule comprises any of the sequences of nucleotides illustrated in any of Sequence ID Nos. 1 to 9.

Whilst the molecules defined herein have been established as being critical for growth and metabolism of *Candida albicans*, for some of the molecules no apparent functionality has been assigned by virtue of the fact that no functionally related sequences in other prokaryotic or eukaryotic organism can be found in respective databases. Thus, advantageously these sequences may be species specific in which case they may be used as selective targets for treatment of diseases mediated by *Candida Albicans* infection. Thus, in one aspect of the invention the nucleic acid molecules preferably comprise the sequences identified in sequence ID Nos. 1, 4, 5 to 9.

In another aspect of the invention the sequences have been arranged functionally and of nucleotides illustrated in Sequence ID Nos. 2 or 3 are preferred and even more preferably in Sequence ID No. 2 and fragments or derivatives of said nucleic acid molecules.

Letters utilised in the sequences according to the invention which are not recognisable as letters of the genetic code signify a position in the nucleic acid sequence where one or more of bases A, G, C or T can occupy the nucleotide position. Representative letters used to identify the range of bases which can be used are as follows:

30

M: A or C
R: A or G
W: A or T
S: C or G
Y: C or T
K: G or T
V: A or C or G

35

H: A or C or T
D: A or G or T
B: C or G or T
N: G or A or T or C

5

In one embodiment of each of the above-identified aspects of the invention the nucleic acid may comprise a mRNA molecule or alternatively a DNA and preferably a cDNA molecule.

10 Also provided by the present invention is a nucleic acid molecule capable of hybridising to the nucleic acid molecules illustrated in any of Figures 1 to 9 under high stringency conditions such as antisense molecule and which conditions are generally
15 known to those of skill in the art.

Stringency of hybridisation as used herein refers to conditions under which polynucleic acids are stable. The stability of hybrids is reflected in the melting temperature (T_m) of the hybrids. T_m can be
20 approximated by the formula:

$$81.5^{\circ}\text{C} + 16.6 (\text{Log}_{10} [\text{Na}^+] + 0.41 (\% \text{G\&C}) - 600 \text{L/L})$$

wherein L is the length of the hybrids in nucleotides.

25 T_m decreases approximately by 1-1.5°C with every 1% decrease in sequence homology.

The term "stringency" refers to the hybridisation conditions wherein a single-stranded nucleic acid joins with a complementary strand when the purine or
30 pyrimidine bases therein pair with their corresponding base by hydrogen bonding. High stringency conditions favour homologous base pairing whereas low stringency conditions disfavour non-homologous base pairing.

"Low stringency" conditions comprise, for
35 example, a temperature of about 37°C or less, a formamide concentration of less than about 50%, and a moderate to low salt (SSC) concentration; or,

alternatively, a temperature of about 50°C or less, and a moderate to high salt (SSPE) concentration, for example 1M NaCl.

"High stringency" conditions comprise, for
5 example, a temperature of about 42°C or less, a formamide concentration of less than about 20%, and a low salt (SSC) concentration; or, alternatively, a temperature of about 65°C, or less, and a low salt (SSPE) concentration. For example, high stringency
10 conditions comprise hybridization in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C (Ausubel, F.M. et al. Current Protocols in Molecular Biology, Vol. I, 1989; Green Inc. New York, at 2.10.3).

15 "SSC" comprises a hybridization and wash solution. A stock 20X SSC solution contains 3M sodium chloride, 0.3M sodium citrate, pH 7.0.

"SSPE" comprises a hybridization and wash solution. A 1X SSPE solution contains 180 mM NaCl,
20 9mM Na₂HPO₄ and 1 mM EDTA, pH 7.4.

The nucleic acid capable of hybridising to nucleic acid molecules according to the invention will generally be at least 70%, preferably at least 80 or 90% and more preferably at least 95% homologous to the
25 nucleotide sequences illustrated in any of Figures 1 to 9.

The DNA molecules according to the invention may, advantageously, be included in a suitable expression vector to express polypeptides encoded therefrom in a
30 suitable host which are critical for growth and survival of *Candida albicans*.

An expression vector according to the invention includes a vector having a nucleic acid according to the invention operably linked to regulatory sequences,
35 such as promoter regions, that are capable of effecting expression of said DNA fragments. The term

"operably linked" refers to a juxtaposition wherein the components described are in a relationship permitting them to function in their intended manner. Such vectors may be transformed into a suitable host cell to provide for expression of a polypeptide according to the invention. Thus, in a further aspect, the invention provides a process for preparing polypeptides according to the invention which comprises cultivating a host cell, transformed, transfected or infected with an expression vector as described above under conditions to provide for expression by the vector of a coding sequence encoding the polypeptides, and recovering the expressed polypeptides.

15 The vectors may be, for example, plasmid, virus or phage vectors provided with an origin of replication, optionally a promoter for the expression of said nucleotide and optionally a regulator of the promoter. The vectors may contain one or more
20 selectable markers, such as, for example, ampicillin resistance.

Regulatory elements required for expression include promoter sequences to bind RNA polymerase and transcription initiation sequences for ribosome
25 binding. For example, a bacterial expression vector may include a promoter such as the lac promoter and for translation initiation the Shine-Dalgarno sequence and the start codon AUG. Similarly, a eukaryotic expression vector may include a heterologous or
30 homologous promoter for RNA polymerase II, a downstream polyadenylation signal, the start codon AUG, and a termination codon for detachment of the ribosome. Such vectors may be obtained commercially or assembled from the sequences described by methods
35 well known in the art.

Polynucleotides according to the invention may be inserted into the vectors described in an antisense

orientation in order to provide for the production of antisense RNA. Antisense RNA or other antisense nucleic acids may be produced by synthetic means.

In accordance with the present invention, a
5 defined nucleic acid includes not only the identical nucleic acid but also any minor base variations including in particular, substitutions in bases which result in a synonymous codon (a different codon specifying the same amino acid residue) due to the
10 degenerate code. The term "nucleic acid sequence" also includes the complementary sequence to any single stranded sequence given regarding base variations.

The present invention also comprises within its scope proteins or polypeptides expressed by the
15 nucleic acid molecules according to the invention or a functional equivalent, derivative or bioprecursor thereof.

The present invention also advantageously provides nucleic acid sequences of at least
20 approximately 10 contiguous nucleotides of a nucleic acid according to the invention and preferably from 10 to approximately 120 nucleotides. In another aspect of the invention, nucleotide acid sequences are provided from 10 to 50 nucleotides. These sequences
25 may, advantageously be used as probes or primers to initiate replication, or the like. Such nucleic acid sequences may be produced according to techniques well known in the art, such as by recombinant or synthetic means. They may also be used in diagnostic kits or
30 the like for detecting the presence of a nucleic acid according to the invention. These tests generally comprise contacting the probe with the sample under hybridising conditions and detecting for the presence of any duplex or triplex formation between the probe
35 and any nucleic acid in the sample.

According to the present invention, these probes may be anchored to a solid support. Preferably, they

- are present on an array so that multiple probes can simultaneously hybridise to a single biological sample. The probes can be spotted onto the array or synthesized *in situ* on the array. See Lockhart et al.,
5 Nature Biotechnology, Vol. 14, December 1996, "Expression monitoring by hybridization to high-density oligonucleotide arrays". A single array can contain more than up to more than a million different probes in discrete locations.
- 10 Advantageously, the nucleic acid sequences, according to the invention may be produced using such recombinant or synthetic means, such as for example using PCR cloning mechanisms which generally involve making a pair of primers, which may be between
15 approximately 10 to 120 nucleotides to a region of the gene which is desired to be cloned, bringing the primers into contact with mRNA, cDNA, or genomic DNA from a cell, performing a polymerase chain reaction under conditions which bring about amplification of
20 the desired region, isolating the amplified region or fragment and recovering the amplified DNA. Generally, such techniques as defined herein are well known in the art, such as described in Sambrook et al (Molecular Cloning: a Laboratory Manual, 1989).
- 25 The nucleic acids or oligonucleotides according to the invention may carry a revealing label. Suitable labels include radioisotopes such as ^{32}P or ^{35}S , enzyme labels or other protein labels such as biotin or fluorescent markers. Such labels may be
30 added to the nucleic acids or oligonucleotides of the invention and may be detected using known techniques *per se*.

The polypeptide or protein according to the invention includes all possible amino acid variants
35 encoded by the nucleic acid molecule according to the invention including a polypeptide encoded by said molecule and having conservative amino acid changes.

Nucleic acids and polypeptides which are particularly preferred are those comprising the sequences of nucleotides provided in SEQ ID NOS: 1-3 and polypeptides provided in SEQ ID NOS: 10-12. However, a particularly preferred nucleic acid comprises the sequences of nucleotides provided as SEQ ID NOS: 2 and/or 3, and their corresponding amino acid sequences identified as SEQ ID NOS: 11 and 12.

With reference to the nucleic acids of this invention and the protein encoded thereby, amino acid sequence SEQ ID NO: 10 is the translation of SEQ ID NO:1; SEQ ID NO:1 is the translation of SEQ ID NO:2; SEQ ID NO:12 is the translation of SEQ ID NO:3; SEQ ID NO: 16 is the translation of SEQ ID NO:4; SEQ ID NO:13 is the translation of SEQ ID NO:5; SEQ ID NO:14 is the translation of SEQ ID NO:6 and SEQ ID NO:15 is the translation of SEQ ID NO:7.

Nucleotide sequences according to the invention are particularly advantageous as selective therapeutic targets for treating *Candida albicans* associated infections. For example, an antisense nucleic acid capable of binding to the nucleic acid sequence illustrated in any of SEQ ID NOS: 1-9 may be used to selectively inhibit expression of the corresponding polypeptides, leading to impaired growth of the *Candida albicans* with reductions of associated illnesses or diseases (see Figure 5).

The nucleic acid molecule or the polypeptide according to the invention may be used as a

medicament, or in the preparation of a medicament, for treating diseases or conditions associated with *Candida albicans* infection.

- Advantageously, the nucleic acid molecule or the
5. polypeptide according to the invention may be provided in a pharmaceutical composition together with a pharmaceutically acceptable carrier, diluent or excipient therefor.

- The present invention is further directed to
- 10 inhibiting expression of nucleic acids according to the invention *in vivo* by the use of antisense technology. Antisense technology can be used to control gene expression through triple-helix formation of antisense DNA or RNA, both of which methods are
- 15 based on binding of a polynucleotide to DNA or RNA. For example, the 5' coding portion or the mature protein sequence, which encodes for the protein of the present invention, is used to design an antisense RNA oligonucleotide of from 10 to 50 base pairs in length.
20. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription (triple-helix - see Lee et al. Nucl. Acids Res., 6:3073 (1979); Cooney et al., Science, 241:456 (1988); and Dervan et al., Science, 251: 1360 (1991), thereby
- 25 preventing transcription and the production of the corresponding protein. The antisense RNA oligonucleotide hybridises to the mRNA *in vivo* and blocks translation of an mRNA molecule into the corresponding protein (antisense - Okano, J.,
- 30 Neurochem., 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988)).

- Antibodies to the protein or polypeptide of the present invention may, advantageously, be prepared by
- 35 techniques which are known in the art. For example, polyclonal antibodies may be prepared by inoculating a host animal, such as a mouse, with the polypeptide

according to the invention or an epitope thereof and recovering immune serum. Monoclonal antibodies may be prepared according to known techniques such as described by Kohler R. and Milstein C., Nature (1975)
5 256, 495-497.

Antibodies according to the invention may also be used in a method of detecting for the presence of a polypeptide according to the invention, which method comprises reacting the antibody with a sample and
10 identifying any protein bound to said antibody. A kit may also be provided for performing said method which comprises an antibody according to the invention and means for reacting the antibody with said sample.

Proteins which interact with the polypeptide of the invention may be identified by investigating
15 protein-protein interactions using the two-hybrid vector system first proposed by Chien et al. (1991).

This technique is based on functional reconstitution *in vivo* of a transcription factor which
20 activates a reporter gene. More particularly the technique comprises providing an appropriate host cell with a DNA construct comprising a reporter gene under the control of a promoter regulated by a transcription factor having a DNA binding domain and an activating
25 domain, expressing in the host cell a first hybrid DNA sequence encoding a first fusion of a fragment or all of a nucleic acid sequence according to the invention and either said DNA binding domain or said activating domain of the transcription factor, expressing in the
30 host at least one second hybrid DNA sequence, such as a library or the like, encoding putative binding proteins to be investigated together with the DNA binding or activating domain of the transcription factor which is not incorporated in the first fusion;
35 detecting any binding of the proteins to be investigated with a protein according to the invention by detecting for the presence of any reporter gene

product in the host cell; optionally isolating second hybrid DNA sequences encoding the binding protein.

An example of such a technique utilises the GAL4 protein in yeast. GAL4 is a transcriptional activator of galactose metabolism in yeast and has a separate domain for binding to activators upstream of the galactose metabolising genes as well as a protein binding domain. Nucleotide vectors may be constructed, one of which comprises the nucleotide residues encoding the DNA binding domain of GAL4. These binding domain residues may be fused to a known protein encoding sequence, such as for example the nucleic acids according to the invention. The other vector comprises the residues encoding the protein binding domain of GAL4. These residues are fused to residues encoding a test protein. Any interaction between polypeptides encoded by the nucleic acid according to the invention and the protein to be tested leads to transcriptional activation of a reporter molecule in a GAL4 transcription deficient yeast cell into which the vectors have been transformed. Preferably, a reporter molecule such as β -galactosidase is activated upon restoration of transcription of the yeast galactose metabolism genes.

Further provided by the present invention is one or more *Candida albicans* cells comprising an induced mutation in the DNA sequence encoding the polypeptide according to the invention.

A further aspect of the invention provides a method of identifying compounds which selectively inhibit or interfere with the expression, the functionality of polypeptides expressed from the nucleotides sequences illustrated in any of SEQ ID NOS: 1-9, or the metabolic pathways in which these polypeptides are involved and which are critical for growth and survival of *Candida albicans*, which method comprises (a) contacting a compound to be tested with

one or more *Candida albicans* cells having a mutation in a nucleic acid molecule according to the invention which mutation results in overexpression or underexpression of said polypeptides in addition to
5 one or more wild type *Candida* cells, (b) monitoring the growth and/or activity of said mutated cell compared to said wild type wherein differential growth or activity of said one or more mutated *Candida* cells provides an indication of selective action of said
10 compound on said polypeptide or another polypeptide in the same or a parallel pathway.

Compounds identifiable or identified using the method according to the invention, may advantageously be used as a medicament, or in the preparation of a
15 medicament to treat diseases or conditions associated with *Candida albicans* infection. These compounds may also advantageously be included in a pharmaceutical composition together with a pharmaceutically acceptable carrier, diluent or excipient therefor.

20 A further aspect of the invention provides a method of identifying DNA sequences from a cell or organism which DNA encodes polypeptides which are critical for growth or survival, which method comprises (a) preparing a cDNA or genomic library from
25 said cell or organism in a suitable expression vector which vector is such that it can either integrate into the genome in said cell or that it permits transcription of antisense RNA from the nucleotide sequences in said cDNA or genomic library, (b)
30 selecting transformants exhibiting impaired growth and determining the nucleotide sequence of the cDNA or genomic sequence from the library included in the vector from said transformant. Preferably, the cell or organism may be any yeast or filamentous fungus,
35 such as, for example, *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe* or *Candida albicans*.

A further aspect of the invention provides a

pharmaceutical composition comprising any of a compound, an antisense molecule or an antibody according to the invention together with a pharmaceutically acceptable carrier, diluent or
5 excipient therefor.

The antisense molecules or indeed the compounds identified as agonists or antagonists of the nucleic acids or polypeptides according to the invention may be used in the form of a pharmaceutical composition,
10 which may be prepared according to procedures well known in the art. Preferred compositions include a pharmaceutically acceptable vehicle or diluent or excipient, such as for example, a physiological saline solution. Other pharmaceutically acceptable carriers
15 including other non-toxic salts, sterile water or the like may also be used. A suitable buffer may also be present allowing the compositions to be lyophilized and stored in sterile conditions prior to reconstitution by the addition of sterile water for
20 subsequent administration. Incorporation of the polypeptides of the invention into a solid or semi-solid biologically compatible matrix may be carried out which can be implanted into tissues requiring treatment.

25 The carrier can also contain other pharmaceutically acceptable excipients for modifying other conditions such as pH, osmolarity, viscosity, sterility, lipophilicity, solubility or the like. Pharmaceutically acceptable excipients which permit
30 sustained or delayed release following administration may also be included.

The polypeptides, the nucleic acid molecules or compounds according to the invention may be administered orally. In this embodiment they may be
35 encapsulated and combined with suitable carriers in solid dosage forms which would be well known to those skilled in the art.

As would be well known to those of skill in the art, the specific dosage regime may be calculated according to the body surface area of the patient or the volume of body space to be occupied, dependent upon the particular route of administration to be used. The amount of the composition actually administered will, however, be determined by a medical practitioner, based on the circumstances pertaining to the disorder to be treated, such as the severity of the symptoms, the composition to be administered, the age, weight, and response of the individual patient and the chosen route of administration.

Example 1

15 Identification of novel drug targets in *C. albicans* by anti-sense and disruptive integration

The principle of the approach is based on the fact that when a particular *C. albicans* mRNA is inhibited by producing the complementary anti-sense RNA, the corresponding protein will decrease. If this protein is critical for growth or survival, the cell producing the anti-sense RNA will grow more slowly or will die.

Since anti-sense inhibition occurs at mRNA level, the gene copy number is irrelevant, thus allowing applications of the strategy even in diploid organisms.

Anti-sense RNA is endogenously produced from an integrative or episomal plasmid with an inducible promoter; induction of the promoter leads to the production of an RNA encoded by the insert of the plasmid. This insert will differ from one plasmid to another in the library. The inserts will be derived from genomic DNA fragments or from cDNA to cover to the extent possible- the entire genome.

The vector is a proprietary vector allowing

integration by homologous recombination at either the homologous insert or promoter sequence in the *Candida* genome. After introducing plasmids from cDNA or genomic libraries into *C. albicans*, transformants are
5 screened for impaired growth after promoter (& thus anti-sense) induction in the presence of lithium acetate. Lithium acetate prolongs the G1 phase and thus allows anti-sense to act during a prolonged period of time during the cell cycle. Transformants
10 which show impaired growth in both induced and non-induced media, thus showing a growth defect due to integrative disruption, are selected as well.

Transformants showing impaired growth are supposed to contain plasmids which produce anti-sense
15 RNA to mRNAs critical for growth or survival. Growth is monitored by measuring growth-curves over a period of time in a device (Bioscreen Analyzer, Labsystems) which allows simultaneous measurement of growth-curves of 200 transformants.

20 Subsequently plasmids can be recovered from the transformants and the sequence of their inserts determined, thus revealing which mRNA they inhibit. In order to be able to recover the genomic or cDNA insert which has integrated into the *Candida* genome, genomic
25 DNA is isolated, cut with an enzyme which cuts only once into the library vector (and estimated approx. every 4096 bp in the genome) and religated. PCR with primers flanking the insert will yield (partial) genomic or cDNA inserts as PCR fragments which can
30 directly be sequenced. This PCR analysis (on ligation reaction) will also show us how many integrations occurred. Alternatively the ligation reaction is transformed to *E. coli* and PCR analysis is performed on colonies or on plasmid DNA derived thereof.

35 This method is employed for a genome-wide search for novel *C. albicans* genes which are important for growth or survival.

Materials & Methods

Construction of pGAL1PNiST-1 (SEQ ID NO:21)

The backbone of the pGAL1PNiST-1 vector
5 (integrative anti-sense *SfiI*-*NotI* vector) is
pGEM11Zf(+) (Promega Inc.). First, the CaMAL2
EcoRI/*SalI* promoter fragment from pDBV50 (D.H. Brown
et al. 1996) was ligated into *EcoRI*/*SalI*-opened
pGEM11Zf(+) resulting in the intermediate construct
10 pGEMMAL2P-1. Into the latter (*MscI*/*CIP*) the CaURA3
selection marker was cloned as a *Eco47III*/*XmnI*
fragment derived from pRM2. The resulting pGEMMAL2P-2
vector was *NotI*/*HindIII* opened in order to accept the
NotI-stuffer-*SfiI* cassette from pPCK1NiSCYCT-1
15 (*EagI*/*HindIII* fragment): pMAL2PNiST-1. Finally, the
plasmid pGAL1PNiST-1 was constructed by exchanging the
SalI/*Ecl136II* MAL2 promoter in pMAL2PNiST-1 by the
XhoI/*SmaI* GAL1 promoter fragment derived from
pRM2GAL1P.

20

Construction of pGAL1PSiST-1 (SEQ ID ON:22)

The vector pGAL1PSiST-1 was created for cloning
the small genomic DNA fragments (flanked by *SfiI*
sites) behind the GAL1 promoter. The only difference
25 with pGAL1PNiST-1 is that the hIFN β (stuffer fragment)
insert fragment in pGAL1PSiST-1 is flanked by two *SfiI*
sites in stead of a *SfiI* and a *NotI* site as in
pGAL1PNiST-1. To construct pGAL1PSiST-1 the *EcoRI*-
HindIII fragment, containing hIFN β flanked by a *SfiI*
30 and a *NotI* site, of pMAL2pHiET-3 (unpublished) was
exchanged by the *EcoRI*-*HindIII* fragment, containing
hIFN β flanked by two *SfiI* sites, from YCp50S-S (an *E.*
coli / *S. cerevisiae* shuttle vector derived from the
plasmid YCp50, which is deposited in the ATCC
35 collection (number 37419; Thrash et al., 1985); an

EcoRI-HindIII fragment, containing the gene *hIFN β* , which is flanked by two *SfiI* sites, was inserted in YCp50, creating YCp50S-S), resulting into plasmid pMAL2PSiST-1. The *MAL2* promoter from pMAL2PSiST-1 (by
5 a *NaeI-balI* digest) was further replaced by the *GAL1* promoter from pGAL1PNiST-1 (via a *XhoI-FSPI* digest), creating the vector pGAL1PSiST-1.

***Candida albicans* genomic library**

10 * Preparation of the genomic DNA fragments

A *Candida albicans* genomic DNA library with small DNA fragments (400 to 1,000 bp) was prepared. Genomic DNA of *Candida albicans* B2630 was isolated following a modified protocol of Blin and Stafford (1976). The
15 quality of the isolated genomic DNA was checked by gel electrophoresis. Undigested DNA was located on the gel above the marker band of 26,282 bp. A little smear, caused by fragmentation of the DNA, was present. To obtain enrichment for genomic DNA fragments of the
20 desired size, the genomic DNA was partially digested. Several restriction enzymes (*AluI*, *HaeIII* and *RsaI*; all creating blunt ends) were tried out. The appropriate digest conditions have been determined by titration of the enzyme. Enrichment of small DNA
25 fragments was obtained with 70 units of *AluI* on 10 μ g of genomic DNA for 20 min. T4 DNA polymerase (Boehringer) and dNTPs (Boehringer) were added to polish the DNA ends. After extraction with phenol-chloroform the digest was size-fractionated on an
30 agarose gel. The genomic DNA fragments with a length of 500 to 1,250 bp were eluted from the gel by centrifugal filtration (Zhu et al., 1985). *SfiI* adaptors (5' GTTGGCCTTTT) or (5' AGGCCAAC) were attached to the DNA ends (blunt) to facilitate cloning
35 of the fragments into the vector. Therefore, a 8-mer and 11-mer oligonucleotide (comprising the *SfiI* site)

were kinased and annealed. After ligation of these adaptors to the DNA fragments a second size-fractionation was performed on an agarose gel. The DNA fragments of 400 to 1150 bp were eluted from the gel by centrifugal filtration.

* *Preparation of the pGAL1PSiST-1 vector fragment*

The small genomic DNA fragments were cloned after the GAL1 promoter in the vector pGAL1PSiST-1. Qiagen-purified pGAL1PSiST-1 plasmid DNA was digested with *SfiI* and the largest vector fragment eluted from the gel by centrifugal filtration (Zhu *et al.*, 1985). Ligation with a control DNA fragment, flanked by *SfiI* sites, was performed as a control. The ligation mix was electroporated to MC1061 *E. coli* cells. Plasmid DNA of 24 clones was analyzed. In all cases the control fragment was inserted in the pGAL1PSiST-1 vector fragment.

* *Upscaling*

All genomic DNA fragments (450 ng) were ligated into the pGAL1PSiST-1 vector (20-ng). After electroporation at 2500V, 40 μ F circa 400,000 clones were obtained. These clones were pooled into three groups and stored as glycerol slants. Also Qiagen-purified DNA was prepared from these clones. A clone analysis showed an average insert length of 600 bp and a percentage of 91 for clones with an insert. The size of the library corresponds to 5 times the diploid genome. The genomic DNA inserts are sense or anti-sense orientated in the vector.

30

***Candida albicans* cDNA library**

Total RNA was extracted from *Candida albicans* B2630 grown on respectively minimal (SD) and rich (YPD) medium as described by Chirgwin *et al.* in Sambrook *et al* 1996. mRNA was prepared from total RNA using the Invitrogen Fast Track procedure.

First strand cDNA is synthesised with the

Superscript Reverse Transcriptase (BRL) and with an oligo dT-NotI Primer adapter. After second strand synthesis, cDNA is polished with Klenow enzyme and purified over a Sephacryl S-400 spun column.

- 5 Phosphorylated *SfiI* adapters are then ligated to the cDNA, followed by digestion with the *NotI* restriction enzyme. The *SfiI*/*NotI* cDNA is then purified and sized on a Biogel column A150M.

First fraction contains approximately 38,720 clones by transformation, the second fraction only 1540 clones. Clone analysis:

Fr. I: 22/24 inserts, 16 ; 1000 bp, 4 ; 2000 bp, average size: 1500 bp.

Fr. II: 9/12 inserts, 3 ; 1000 bp, average size: 960

- 15 bp cDNA was ligated in a *NotI*/*SfiI* opened pGAL1PNiST-1 vector (anti-sense)

Candida transformation

The host strain used for transformation is a *C. albicans* *ura3* mutant, CAI-4, which contains a deletion in orotidine-5-phosphate decarboxylase and was obtained from William Fonzi, Georgetown University (Fonzi and Irwin). CAI-4 was transformed with the above described cDNA library or genomic library using the *Pichia* spheroplast module (Invitrogen). Resulting transformants were plated on minimal medium supplemented with glucose (SD, 0.67% or 1.34% Yeast Nitrogen base w/o amino acids + 2% glucose) plates and incubated for 2-3 days at 30°C.

30

Screening for mutants

Starter cultures were set up by inoculating each colony in 1 ml SD medium and incubating overnight at 30°C and 300 rpm. Cell densities were determined using a Coulter counter (Coulter Z1; Coulter electronics limited). 250.000 cells/ml were inoculated in 1 ml SD

35

medium and cultures were incubated for 24 hours at 30°C and 300 rpm. Cultures were washed in minimal medium without glucose (S) and the pellet resuspended in 650 µl S medium. 8 µl of this culture is used for
5 inoculating 400 µl cultures in a Honeywell-100 plate (Bioscreen analyzer; Labsystems). Each transformant was grown during three days in S medium containing LiAc; pH 6.0, with 2% glucose/2% maltose or 2% galactose/2% maltose respectively while shaking every
10 3 minutes for 20 seconds. Optical densities were measured every hour during three consecutive days and growth curves were generated (Bioscreen analyzer; Labsystems).

Growth curves of transformants grown in
15 respectively anti-sense non-inducing (glucose/maltose) and inducing (galactose/maltose) medium are compared and those transformants showing impaired growth upon anti-sense induction are selected for further analysis. Transformants showing impaired growth by
20 virtue of integration into a critical gene are also selected.

Isolation of genomic or cDNA inserts

Putatively interesting transformants are grown in
25 1.5 ml SD overnight and genomic DNA is isolated using the Nucleon MI Yeast kit (Clontech). Concentration of genomic DNA is estimated by analyzing a sample on an agarose gel.

20 ng of genomic DNA is digested for three hours
30 with an enzyme that cuts uniquely in the library vector (SacI for the genomic library; PstI for the cDNA library) and treated with RNase. Samples are phenol/chloroform extracted and precipitated using NaOAc/ethanol.

35 The resulting pellet is resuspended in 500 µl ligation mixture (1 x ligation buffer and 4 units of

T4 DNA ligase; both from Boehringer) and incubated overnight at 16°C.

After denaturation (20 min 65°C), purification (phenol/chloroform extraction) and precipitation (NaOAc/ethanol) the pellet is resuspended in 10 µl MilliQ (Millipore) water.

PCR analysis

Inverse PCR is performed on 1 µl of the precipitated ligation reaction using library vector specific primers (oligo23 5' TGC-AGC-TCG-ACC-TCG-ACT-G 3' SEQ ID NO: 17 and oligo25 5' GCG-TGA-ATG-TAA-GCG-TGA-C 3' SEQ ID NO: 18 for the genomic library; 3pGALNistPCR primer: 5'TGAGCAGCTCGCCGTCGCGC 3' SEQ ID NO: 19 and 5pGALNistPCR primer: 5'GAGTTATACCCTGCAGCTCGAC 3' SEQ ID NO: 20 for the cDNA library; both from Eurogentec) for 30 cycles each consisting of (a) 1 min at 95 °C, (b) 1 min at 57 °C, and (c) 3 min at 72 °C. In the reaction mixture 2.5 units of Taq polymerase (Boehringer) with TaqStart antibody (Clontech) (1:1) were used, and the final concentrations were 0.2 µM of each primer, 3 mM MgCl₂ (Perkin Elmer Cetus) and 200 µM dNTPs (Perkin Elmer Cetus). PCR was performed in a Robocycler (Stratagene).

Sequence determination

Resulting PCR products were purified using PCR purification kit (Qiagen) and were quantified by comparison of band intensity on EtBr stained agarose gel with the intensity of DNA marker bands. The amount of PCR product (expressed in ng) used in the sequencing reaction is calculated as the length of the PCR product in basepairs divided by 10. Sequencing reactions were performed using the ABI Prism BigDye Terminator Cycle Sequencing Ready Reaction Kit

according to the instructions of the manufacturer (PE Applied Biosystems, Foster City, CA) except for the following modifications.

The total reaction volume was reduced to 15 μ l.

5 Reaction volume of individual reagents were changed accordingly. 6.0 μ l Terminator Ready Reaction Mix was replaced by a mixture of 3.0 μ l Terminator Ready Reaction Mix + 3.0 μ l Half Term (GENPAK Limited, Brighton, UK). After cycle sequencing, reaction

10 mixtures were purified over Sephadex G50 columns prepared on Multiscreen HV opaque microtiter plates (Millipore, Molsheim, Fr) and were dried in a speedVac. Reaction products were resuspended in 3 μ l loading buffer. Following denaturation for 2 min at

15 95°C, 1 μ l of sample was applied on a 5% Long Ranger Gel (36 cm well-to-read) prepared from Singel Packs according to the supplier's instructions (FMC BioProducts, Rockland, ME). Samples were run for 7 hours 2X run on a ABI 377XL DNA sequencer. Data

20 collection version 2.0 and Sequence analysis version 3.0 (for basecalling) software packages are from PE Applied Biosystems. Resulting sequence text files were copied onto a server for further analysis.

25 **Sequence analysis**

Nucleotide sequences were imported in the VectorNTI software package (InforMax Inc, North Bethesda, MD, USA), and the vector and insert regions of the sequences were identified. Sequence similarity

30 searches against public and commercial sequence databases were performed with the BLAST software package (Altschul et al., 1990) version 1.4. Both the original nucleotide sequence and the six-frame conceptual translations of the insert region were used

35 as query sequences. The used public databases were the EMBL nucleotide sequence database (Stoesser et al.,

1998), the SWISS-PROT protein sequence database and its supplement TrEMBL (Bairoch and Apweiler, 1998), and the ALCES *Candida albicans* sequence database (Stanford University, University of Minnesota). The commercial sequence databases used were the LifeSeq[®] human and PathoSeq[®] microbial genomic databases (Incyte Pharmaceuticals Inc., Palo Alto, CA, USA), and the GENESEQ patent sequence database (Derwent, London, UK). Three major results were obtained on the basis of the sequence similarity searches: function, novelty, and specificity. A putative function was deduced on the basis of the similarity with sequences with a known function, the novelty was based on the absence or presence of the sequences in public databases, and the specificity was based on the similarity with vertebrate homologues.

Methods

Blastx of the nucleic acid sequences against the appropriate protein databases: Swiss-Prot for clones of which the complete sequence is present in the public domain, and paorfp (PathoSeq[™]) for clones of which the complete sequences is not present in the public domain.

The protein to which the translated nucleic acid sequence corresponds to is used as a starting point. The differences between this protein and our translated nucleic acid sequences are marked with a double line and annotated above the protein sequence.

The following symbols are used:

A one-letter amino acid code or the ambiguity code X is used if our translated nucleic acid sequence has another amino acid on a certain position,

The stop codon sign * is used if our translated nucleic acid sequence has a stop codon on a certain position,

The letters fs (frame shift) are used if a frame

shift occurs in our translated nucleic acid sequence,
and another reading frame is used,

The words ambiguity or ambiguities are used if a
part of our translated nucleic acid sequence is
5 present in the proteins, but not visible in the
alignments of the blast results,

The phrase "missing sequence" is used if the
translated nucleic acid sequence does not comprise
that part of the protein.

10 Blastx: compares the six-frame conceptual
translation products of a nucleotide query sequence
(both strands) against a protein sequence database.

Antisense Experiments

15

Clones 383c_cp, 392c_cp, 417c_cpG2L, 325c_af,
322c_cp, 26g3, and 409c_cp were transformed with
plasmid pGAL1PSiST-1 containing the galactose
inducible promotor/expression cassette. The plasmid
20 was modified to include antisense molecules capable of
binding to RNA expressed from DNA corresponding to SEQ
ID NOs: 1-2 and SEQ ID NOs 4-8 natively present the
Candida albicans cell. Following transformation the
cells were grown in the presence of glucose or
25 galactose to determine the effect of the induced
antisense molecule on protein expression and on
Candida albicans cell growth. An exemplary growth
curve is provided in Figure 5. All cells transformed
with the antisense molecules demonstrated significant
30 growth reduction.

Gene Knock-outs

To verify that the growth effect was due to the
interference with the identified gene and to support
35 the specificity of the antisense effect, single and
double allele knock-outs were made in the identified
genes using the URA-blaster method (Fonzi and Irwin

1993). Figure 4 provides an example of a growth curve for a double allele knock-out as compared to the wild-type strain CAI4. Growth curves are shown as the measurement of optical density (Y-axis) over time in hours (X-axis). The optical density is a measure for the number of cells present and this a measure for growth. A reduction in growth was observed for this double knock-out as compared to the wild-type strain.

10 **Screening for compounds modulating expression of**
15 **polypeptides critical for growth and survival of *C.***
albicans

The method proposed is based on observations (Sandbaken et al., 1990; Hinnebusch and Liebman 1991; Ribogene PCT WO 95/11969, 1995) suggesting that underexpression or overexpression of any component of a process (e.g. translation) could lead to altered sensitivity to an inhibitor of a relevant step in that process. Such an inhibitor should be more potent against a cell limited by a deficiency in the macromolecule catalyzing that step and/or less potent against a cell containing an excess of that macromolecule, as compared to the wild type (WT) cell.

Mutant yeast strains, for example, have shown that some steps of translation are sensitive to the stoichiometry of macromolecules involved. (Sandbaken et al. 1996). Such strains are more sensitive to compounds, which specifically perturb translation (by acting on a component that participates in translation) but are equally sensitive to compounds with other mechanisms of action.

This method thus not only provides a means to identify whether a test compound perturbs a certain process but also an indication of the site at which it exerts its effect. The component which is present in altered form or amount in a cell whose growth is

affected by a test compound is potentially the site of action of the test compound.

The assay involves measurement of growth of an isogenic strain which has been modified only in a certain specific allele, relative to a wild type (WT) *C. albicans* strain, in the presence of R-compounds. Strains can be ones in which the expression of a specific essential protein is impaired upon induction of anti-sense or strains which carry disruptions in an essential gene. An *in silico* approach to finding novel essential genes in *C. albicans* will be performed. A number of essential genes identified in this way will be disrupted (in one allele) and the resulting strains can be used for comparative growth screening.

Assay for High Throughput screening for drugs

35 μ l minimal medium (S medium + 2% galactose + 2% maltose) is transferred in a transparent flat-bottomed 96 well plate using an automated pipetting system (Multidrop, Labsystems). A 96-channel pipettor (Hydra, Robbins Scientific) transfers 2.5 μ l of R-compound at 10^{-3} M in DMSO from a stock plate into the assay plate.

25 The selected *C. albicans* strains (mutant and parent (CAI-4) strain) are stored as glycerol stocks (15%) at -70°C . The strains are streaked out on selective plates (SD medium) and incubated for two days at 30°C . For the parent strain, CAI-4, the medium is always supplemented with 20 $\mu\text{g/ml}$ uridine. A single colony is scooped up and resuspended in 1 ml minimal medium (S medium + 2% galactose + 2% maltose). Cells are incubated at 30°C for 8 hours while shaking at 250 rpm. A 10 ml culture is inoculated at 250,000 cells/ml. Cultures are incubated at 30°C for 24 hours while shaking at 250 rpm. Cells are counted in Coulter counter and the final culture (S medium + 2%

galactose + 2% maltose) is inoculated at 20,000 to 50,000 cells/ml. Cultures are grown at 30°C while shaking at 250 rpm until a final OD of 0.24 (+/- 0.04) 600nm is reached.

- 5 200 µl of this yeast suspension is added to all wells of MW96 plates containing R-compounds in a 450 (or 250) µl total volume. MW96 plates are incubated (static) at 30°C for 48 hours.

Optical densities are measured after 48 hours.

- 10 Test growth is expressed as a percentage of positive control growth for both mutant (x) and wild type (y) strains. The ratio (x/y) of these derived variables is calculated.

References

- Altschul, S.F., W. Gish, W. Miller, E.W. Myers, and
D.J. Lipman, 1990, Basic Local Alignment Search Tool,
5 J. Mol. Biol. 215: 403-410.
- Thrash C., Bankier A. T., Barrell B. G. and Sternglanz
R. (1985) *Proc. Natl. Acad. Sci. USA* **82**: 4374-4378.
- 10 Blin N. and Stafford D. W. (1976) *Nucleic Acids Res.*
3: 2303-2308.
- Zhu J., Kempnaers W., Van Der Straeten D., Contreras
R., and Fiers W. (1985) *Bio/Technology* **3**: 1014-1016.
- 15 Sambrook J., Fritsch E.F. and Maniatis T. (1989)
Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold
Spring Harbor Laboratory, Cold Spring Harbor, NY.
- 20 ~~Fonzi W. and Irwin H. (1993) *Genetics* **134**:717-728.~~
Altschul S.F, Gish W., Miller W., Myers E.W., Lipman
D.J. (1990) *J. Mol. Biol.* **215**(3):403-410.
- Bairoch A. and Apweiler R. (1998) *Nucleic Acids Res.*
25 **26**(1):38-42.
- Stoesser G., Moseley M.A., Sleep J., McGowran M.,
Garcia-Pastor M., Sterk P. (1998) *Nucleic Acids Res.*
26(1):8-15.
- 30 Chien et al., (1991) *Proc. Natl. Acad. Sci USA* **88**,
9578-9582.
- Sandbaken M.G. Lupisella J.A., DiDomenico B. and
35 Chakraborty K., *J. Biol. Chem.* **265**:15838-15844, 1990.
- Hinnebusch, A.G. and Liebman S.W., in: The Molecular

Biology of the Yeast *Saccharomyces*, Broach J.R.,
Pringle J.R. and Jones E.W., eds., CSH Laboratory
Press; NY 1991.

- 5 Patent application RiboGene Inc., PCT WO 95./11969,
1995.

Brown, D.H. et al., Molecular General Genetics 251 (1)
75-80, 1996.

10

Altschul, S.F., W. Gish, W. Miller, E.W. Myers, and
D.J. Lipman. 1990. Basic local alignment search tool.
J. Mol. Biol. 215: 403-410.

- 15 Arndt, G.M., D. Atkins, M. Patrikakis, and J.G.
Izant 1995. Gene regulation by antisense RNA in the
fission yeast *Schizosaccharomyces pombe*. Mol. Gen.
Genet. 248: 293-300.

- 20 Atkins, D., and W.L. Gerlach. 1994. Artificial
ribozyme and antisense gene expression in *S.*
cerevisiae. Antisense research and development 4:109-
117.

- 25 Atkins, D., G.M. Arndt, and J.G. Izant 1994.
Antisense gene expression in yeast. Biol. Chem. Hoppe-
Seyler. 375:721-729.

- Atkins, D., M. Patrikakis, J.G. Izant 1995. The
30 *ade6* gene of the fission yeast as a target for
antisense and ribozyme RNA-mediated suppression.
Antisense Research & Development. 5(4):295-305.

- Bairoch, A., and R. Apweiler. 1998. The SWISS-PROT
35 protein sequence data bank and its supplement
TrEMBL in 1998. Nucleic Acids Res. 26: 38-42.
Baudin, A., O. Ozier-Kalogeropoulos, A. Denouel, F.

- Lacroute, C. Cullin. 1993. A simple and efficient method for direct gene deletion in *Saccharomyces cerevisiae*. *Nucleic Acids Research*. 21(14):3329-30.
- 5 Blin, N., and D.W. Stafford. 1976. *Nucleic Acids Res.* 3: 2303-2308.
- Dujon, B. 1998. European Functional Analysis Network (EUROFAN) and the functional analysis of the
- 10 *Saccharomyces cerevisiae* genome. *Electrophoresis*. 19:617-624.
- Del Rosario, M., J.C. Stephans, J. Zakel, J. Escobedo, and K. Giese. 1996. Positive selection system to
- 15 screen for inhibitors of human immunodeficiency virus-1 transcription. *Nature Biotechnology*. 14:1592-1596.
- Fairhead, C., A. Thierry, F. Denis, M. Eck, and B. Dujon. 1998. A Mass-murder of ORFs from three regions
- 20 of chromosome XI from *Saccharomyces cerevisiae*. *Gene*. 223:33-46.
- Ferbeyre, G., J. Bratty, H. Chen, R. Cedergren. 1996. Cell cycle arrest promotes trans-hammerhead ribozyme
- 25 action in yeast. *Journal of Biological Chemistry*. 271(32):19318-23.
- Fleischmann, R.D., M.D. Adams, O. White, R.A. Clayton, E.F. Kirkness, A.R. Kerlavage, C.J. Bult, J.F. Tomb,
- 30 B.A. Dougherty, J.M. Merrick. 1995. Whole-genome random sequencing and assembly of *Haemophilus influenzae*. *Science*. 269: 496-512.
- Fonzi, W.A., and M.Y. Irwin. 1993. Isogenic strain
- 35 construction and gene mapping in *Candida albicans*. *Genetics* 134:717-728.
- Fraser, C.M., J.D. Gocayne, O. White, M.D. Adams, R.A.

Clayton, R.D. Fleischmann, C.J. Bult, A.R. Kervalage,
G. Sutton, J.M. Kelley. 1995. The minimal gene
complement of *Mycoplasma genitalium*. *Science* 270:397-
403.

5

Giese, K. 1997. Method and construct for screening for
inhibitors of transcriptional activation.
International patent application WO 97/10360.

- 10 Hahn, S. Guarente L. 1988. Yeast HAP2 and HAP3:
transcriptional activators in a heteromeric complex.
Science. 240(4850):317-21

- Heid, C.A., J. Stevens, K.J. Livak, and P.M. Williams.
15 1996. Real time quantitative PCR. *Genome Methods*
6:986-994.

- Jayaram, M., A. Sutton, J.R. Broach. 1985. Properties
of REP3: a cis-acting locus required for stable
20 propagation of the *Saccharomyces cerevisiae* plasmid 2
microns circle. *Molecular & Cellular Biology*.
5(9):2466-75.

- Jeong, S.W., W.H. Lang, R.H. Reeder. 1996. The yeast
25 transcription terminator for RNA polymerase I is
designed to prevent polymerase slippage. *Journal of*
Biological Chemistry. 271(27):16104-10.

- Johnson, F.B., D.A. Sinclair, and L. Guarente. 1999.
30 *Molecular Biology of aging*. *Cell*. 96:291-302.

- Leuker, C.E., A. Sonneborn, S. Delbruck, J.F. Ernst.
1997. Sequence and promoter regulation of the PCK1
gene encoding phosphoenolpyruvate carboxykinase of the
35 fungal pathogen *Candida albicans*. *Gene*. 192(2):235-
40.

Lie, Y.S., and C.J. Petropoulos. 1998. Advances in

quantitative PCR technology: 5' nuclease assays.
Current Opinion in Biotechnology 9:43-48.

- 5 Livak, K.J., S.J. Flood, J. Marmaro, W. Giusti, K. Deetz. 1995. Oligonucleotides with fluorescent dyes at opposite ends provide a quenched probe system useful for detecting PCR product and nucleic acid hybridization. *Genome Research*. 4(6):357-62.
- 10 Mandart, E. 1998. Effects of mutations in the *Saccharomyces cerevisiae* RNA14 gene on the abundance and polyadenylation of its transcripts. *Mol. Gen. Genet.* 258:16-25.
- 15 McMahon, S.B., H.A. Van Buskirk, K.A. Dugan, T.D. Copeland, and M.D. Cole. 1998. The novel ATM-related protein TRRAP is an essential cofactor for the c-Myc and E2F oncoproteins. *Cell*. 94:363-74.
- 20 Murray, J.A.H., M. Scarpa, N. Rossi, G. Cesareni. 1987. Antagonistic controls regulate copy number of the yeast 2 μ plasmid. *EMBO J.* 6:4205-4212.
- 25 Nasr, F., A. Bécam, P.P. Slonimski, and C.J. Herbert. 1994. YBR1012, an essential gene from *S. cerevisiae*: construction of an RNA-antisense conditional allele and isolation of a multicopy suppressor. *CR Acad. Sci. Paris*. 317:607-613
- 30 Nasr, F., A. Bécam, S.C. Brown, D. De Nay. P.P. Slonimski, and C.J. Herbert. 1995. Artificial antisense regulation of YBR1012 an essential gene from *S. cerevisiae* which is important for progression through G1/S. *Mol. Gen. Genet.* 249:51-57.
- 35 Nomura, T., N. Fujita, A. Ishihama. 1985. Promoter selectivity of *E. coli* RNA polymerase: analysis of the

promoter system of convergently-transcribed dnaQ-rnh genes. Nucleic Acids Research. 13(21):7647-61.

Orlando, C., P. Pinzani, and M. Pazzagli. 1998.

- 5 Developments in quantitative PCR. Clin. Chem. Lab. Med. 36(5):255-269.

Pla, J., C. Gil, F. Monteoliva, M. Sanchez, and C. Nombela. 1996. Understanding Candida albicans at the
10 molecular level. Yeast. 12:1677-1702.

Reeder, R.H. and W.H. Lang. 1997. Terminating transcription in eukaryotes: lessons learned from RNA polymerase I. Trends in Biochemical Sciences.

- 15 22(12):473-7, 1997

Sambrook, J., E.F. Fritsch, and T. Maniatis. 1989. Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.

20

Sinclair, D.A., L. Guarente. 1997. Extrachromosomal rDNA circles--a cause of aging in yeast. Cell. 91(7):1033-42.

- 25 Smith, V., D. Botstein, and P. O. Brown. 1995. Genetic footprinting: A genomic strategy for determining a gene's function given its sequence. Proc. Natl. Acad. Sci. USA. 92:6479-6483.

- 30 Stoesser, G., Moseley M.A., Sleep J., McGowran M., Garcia-Pastor M., Sterk P. 1998. Nucleic Acids Res. 26(1):8-15.

- Thompson-Jager, S. Domdey H. 1990. The intron of the
35 yeast actin gene contains the promoter for an antisense RNA. Current Genetics. 17(3):269-73.
Thompson, J.R., E. Register, J. Curotto, M. Kurtz, and

- R. Kelly. 1998. An improved protocol for the preparation of yeast cells for transformation by electroporation. *Yeast*. 14:565-571.
- 5 Thrash, C., A.T. Bankier, B.G. Barrell, and R. Sternglanz. 1985. *Proc. Natl. Acad. Sci. USA* 82: 4374-4378.
- 10 Van Duin, M., J. van Den Tol, J.H. Hoeijmakers, D. Bootsma, I.P. Rupp, P. Reynolds, L. Prakash, and S. Prakash. 1989. Conserved pattern of antisense overlapping transcription in the homologous human ERCC-1 and yeast RAD10 DNA repair gene regions. *Molecular & Cellular Biology*. 9(4):1794-8.
- 15 Wach, A., A. Brachat, R. Pohlmann, P. Philippsen. 1994. New heterologous modules for classical or PCR-based gene disruptions in *Saccharomyces cerevisiae*. *Yeast*. 10(13):1793-808.
- 20 Wilson, R.B., D. Davis, A.P. Mitchell. 1999. Rapid hypothesis testing with *Candida albicans* through gene disruption with short homology regions. *Journal of Bacteriology*. 181(6):1868-74.
- 25 Zhu, J., W. Kempenaers, D. Van der Straeten, R. Contreras, and W. Fiers. 1985. A method for fast and pure DNA elution from agarose gels by centrifugal filtration. *Bio/Technology*. 3: 1014-1016.
- 30

Table 1

Seq ID No.	Clone	Function
1	382c_cp	-
2	392c_cp	TUF1

3	-	RAD53
4	417c_cpG2L	-
5	325c_af	-
6	322c_cp ¹	-
7	26g3	-
8	409c_cp	-
9	382c_cpG1L2	-
10	382c_cp (prt)	-
11	392c_cp (prt)	TUF1
12		RAD53
13	325c_af (prt) ²	-
14	322c_cp (prt) ²	-
15	26g3 (prt)	-
16	417c_cp 92L (prt)	-

1. 322c-cp is a member of the UPF0057 protein family. It contains potential transmembrane regions (6-23aa; 30-53aa) and could be low temperature or salt-stress inducible.
2. 325c-af shows similarity to IMP4 yeast and related proteins and it might be involved in rRNA processing in *Candida albicans* in a similar way to IMP4.

What is claimed is:

1. A nucleic acid molecule encoding a polypeptide
which is critical for survival and growth of the yeast
5 *Candida albicans* and which nucleic acid molecule is
selected from the group consisting of SEQ ID NO: 1 to
SEQ ID NO: 9.
2. A nucleic acid molecule encoding a polypeptide
10 which is critical for survival and growth of the yeast
Candida albicans and which nucleic acid molecule is
selected from the group consisting of SEQ ID NO: 1 to
SEQ ID NO: 3.
- 15 3. A nucleic acid molecule encoding a polypeptide
which is critical for survival and growth of the yeast
Candida albicans and which nucleic acid molecule is
selected from SEQ. ID NO: 1 or SEQ. ID. NO: 2 and
fragments or derivatives of said nucleic acid
20 molecules.
4. The nucleic acid molecule of Claim 1 which is mRNA.
5. The nucleic acid molecule of Claim 1 which is DNA.
- 25 6. The nucleic acid molecule according to claim 5
which is cDNA.
7. A first nucleic acid molecule capable of
30 hybridising to a second nucleic acid molecule encoding
a polypeptide which is critical for survival and
growth of the yeast *Candida albicans* and which second
nucleic acid molecule is selected from the group
consisting of SEQ ID NO: 1 to SEQ ID NO: 9 wherein
35 hybridisation occurs under high stringency conditions.
8. An antisense molecule comprising a nucleic acid

molecule capable of hybridising to a nucleic acid molecule encoding a polypeptide which is critical for survival and growth of the yeast *Candida albicans* wherein the nucleic acid molecule is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 9.

9. Cells containing a nucleic acid molecule encoding a polypeptide which is critical for survival and growth of the yeast *Candida albicans* and which nucleic acid molecule is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 9, wherein said cells are bacterial or eukaryotic.

10. A polypeptide encoded by a nucleic acid molecule which is critical for survival and growth of the yeast *Candida albicans* and which nucleic acid molecule is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 9.

11. A polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 10 to SEQ ID NO: 16.

12. A recombinant DNA construct comprising a nucleic acid molecule encoding a polypeptide which is critical for survival and growth of the yeast *Candida albicans* and which nucleic acid molecule is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 9, wherein the nucleic acid molecule is DNA.

13. A recombinant DNA construct comprising a first nucleic acid molecule wherein the first nucleic acid molecule or a molecule complementary to the first nucleic acid molecule is capable of hybridizing a second nucleic acid molecule encoding a polypeptide which is critical for survival and growth of the yeast *Candida albicans* and which second nucleic acid

molecule is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 9 wherein the first nucleic acid molecule is inserted in the antisense orientation.

5

14. A recombinant DNA construct according to claim 12 wherein said recombinant DNA construct is an expression vector.

10 15. The construct according to claim 14 which comprises an inducible promoter.

16. The construct according to claim 14 which comprises a sequence encoding a reporter molecule.

15

17. Cells containing a recombinant DNA construct comprising a nucleic acid molecule encoding a polypeptide which is critical for survival and growth of the yeast *Candida albicans* and which nucleic acid molecule is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 9, wherein said cells are bacterial or eukaryotic.

20

18. The nucleic acid molecule according to claim 1 in a pharmaceutically acceptable carrier, diluent, excipient or buffer.

25

19. A method for treating a *Candida albicans*-associated disease comprising the step of:

30 administering a composition of matter comprising an antisense nucleic acid molecule capable of binding to a *Candida albicans*-originating nucleic acid sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:9.

35

20. A pharmaceutical composition comprising a polypeptide according to claim 10 together with a

pharmaceutically acceptable carrier diluent or excipient therefor.

21. A *Candida albicans* cell comprising an induced
5 mutation in the DNA sequence encoding the polypeptide according to claim 10.

22. A method of identifying compounds which
selectively modulate expression or functionality of
10 polypeptides or metabolic pathways in which these polypeptides are involved and which are crucial for growth and survival of *Candida albicans*, which method comprises:

- 15 (a) contacting a compound to be tested with one or more *Candida albicans* cells having a mutation in a nucleic acid molecule according to claim 1 which mutation results in overexpression or underexpression of said polypeptides in addition to contacting one
20 or more wild type *Candida albicans* cells with said compound; and
- (b) monitoring the growth and/or activity of
said mutated cell compared to said wild
25 type; wherein differential growth or activity of said one or more mutated *Candida* cells is indicative of selective action of said compound on a polypeptide or another polypeptide in the same or a parallel
30 pathway.

23. A compound identifiable according to the method of claim 22.

24. A pharmaceutical composition comprising a
35 compound according to claim 23 together with a pharmaceutically acceptable carrier, diluent or excipient therefor.

25. A method of identifying DNA sequences from a cell or organism which DNA encodes polypeptides which are critical for growth or survival of said cell or

5 organism, which method comprises:

- 10 (a) preparing a cDNA or genomic library from said cell or organism in a suitable expression vector which vector is such that it can either integrate into the genome in said cell or that it permits transcription of antisense RNA from the nucleotide sequences in said cDNA or genomic library; and
- 15 (b) selecting transformants exhibiting impaired growth and determining the nucleotide sequence of the cDNA or genomic sequence from the library included in the vector from said transformant.

-----20 26. The method according to claim 25 wherein said cell or organism is a yeast or filamentous fungus.

27. The method according to claim 25 wherein said cell or organism is selected from the group consisting
25 of *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe* or *Candida albicans*.

28. An antibody capable of binding to a polypeptide encoded by a nucleic acid molecule which is critical
30 for survival and growth of the yeast *Candida albicans* and which nucleic acid molecule is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 9.

29. An oligonucleotide comprising a fragment of from
35 10 to 120 contiguous nucleotides of a nucleic acid molecule which is critical for survival and growth of the yeast *Candida albicans* and which nucleic acid

molecule is selected from the group consisting of SEQ
ID NO: 1 to SEQ ID NO: 9.

30. The oligonucleotide according to claim 29
5 comprising a fragment of from 10 to 50 contiguous
nucleotides.

ABSTRACT**DRUG TARGETS IN *CANDIDA ALBICANS***

5

The present invention is concerned with the identification of genes or functional fragments thereof from *Candida albicans* which are critical for growth and cell division and which genes may be used
10 as selective drug targets to treat *Candida albicans* associated infections. Novel nucleic acid sequences from *Candida albicans* are also provided and which encode the polypeptides which are critical for growth of *Candida albicans*. Methods for the identification of
15 anti-fungal compounds which inhibit fungal or yeast growth are also contemplated.

FIGURE 1




FIGURE 2




FIGURE 3





FIGURE 4



Please type a plus sign (+) inside this box → ☐

PTO/SB/01 (12-97)

Approved for use through 9/30/00. OMB 0651-0032
Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION FOR UTILITY OR DESIGN PATENT APPLICATION (37 CFR 1.63) <input checked="" type="checkbox"/> Declaration Submitted with Initial Filing OR <input type="checkbox"/> Declaration Submitted after Initial Filing (surcharge (37 CFR 1.16 (e)) required)	Attorney Docket Number	JAB 1430-PCT-USA
	First Named Inventor	Contreras, Roland Henri
	COMPLETE IF KNOWN	
	Application Number	/
	Filing Date	
	Group Art Unit	
	Examiner Name	

As a below named inventor, I hereby declare that:

My residence, post office address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

DRUG TARGETS IN CANDIDA ALBICANS

the specification of which (Title of the Invention)

☐ is attached hereto
OR
☐ was filed on (MM/DD/YYYY) **12/06/1999** as United States Application Number or PCT International Application Number **PCT/EP99/09833** and was amended on (MM/DD/YYYY) (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment specifically referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached?	
				YES	NO
98204122.0	EP	12/04/1998	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>

☐ Additional foreign application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto:

I hereby claim the benefit under 35 U.S.C. 119(e) of any United States provisional application(s) listed below.

Application Number(s)	Filing Date (MM/DD/YYYY)

☐ Additional provisional application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto.

[Page 1 of 2]

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

Please type a plus sign (+) inside this box → ☐

PTO/SB/01 (12-97)

Approved for use through 9/30/00. OMB 0651-0032
Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION — Utility or Design Patent Application

I hereby claim the benefit under 35 U.S.C. 120 of any United States application(s), or 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of 35 U.S.C. 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Parent Application or PCT Parent Number	Parent Filing Date (MM/DD/YYYY)	Parent Patent Number (if applicable)

☐ Additional U.S. or PCT international application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto.

As a named inventor, I hereby appoint the following registered practitioner(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith: ☐ Customer Number OR ☐ Registered practitioner(s) name/registration number listed below

Place Customer
Number Bar Code
Label here

Name	Registration Number	Name	Registration Number
Steven P. Berman	24,772	Michael Stark	32,495
Andrea L. Colby	30,194	Ellen C. Coletti	34,140
		Mary A. Appollina	34,087

☐ Additional registered practitioner(s) named on supplemental Registered Practitioner Information sheet PTO/SB/02C attached hereto.

Direct all correspondence to: ☐ Customer Number OR ☒ Correspondence address below

Name	Philip S. Johnson				
Address	Johnson & Johnson				
Address	One Johnson & Johnson Plaza				
City	New Brunswick	State	NJ	ZIP	08933-7003
Country	USA	Telephone	(732) 524-2359	Fax	(732) 524-2808

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Name of Sole or First Inventor:		<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name (first and middle (if any))		Family Name or Surname			
Roland Henri		Contreras			
Inventor's Signature				Date	03/22/01
Residence: City	9000 Gent	State		Country	Belgium
Post Office Address	c/o University of Gent, K.L. Ledeganckstraat 35, B-9000 Gent, Belgium				
Post Office Address					
City	Schelderoode/Merelbeke	State		ZIP	9820
Country	Belgium				

☐ Additional inventors are being named on the _____ supplemental Additional Inventor(s) sheet(s) PTO/SB/02A attached hereto

Please type a plus sign (+) inside this box → ☐

PTO/SB/02A (3-97)
 Approved for use through 9/30/98. OMB 0651-0032
 Patent and Trademark Office, U.S. DEPARTMENT OF COMMERCE
 Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION
ADDITIONAL INVENTOR(S)
Supplemental Sheet
 Page 1 of 2

Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name (first and middle (if any))		Family Name or Surname	
Bart Jozef Maria		Nelissen	
Inventor's Signature	Date		03/20/01
Residence: City	Beerse	State	Country Belgium Citizenship BE
Post Office Address	c/o Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse, Belgium		
Post Office Address			
City	Meerhout	State	ZIP 2450 Country Belgium
Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name (first and middle (if any))		Family Name or Surname	
Marianne Denise		De Backer	
Inventor's Signature	Date		
Residence: City	Beerse	State	Country Belgium Citizenship BE
Post Office Address	c/o Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse, Belgium		
Post Office Address			
City	San Diego	State CA	ZIP 92122 Country USA
Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name (first and middle (if any))		Family Name or Surname	
Walter Herman Maria Louis		Luyten	
Inventor's Signature	Date		03/20/01
Residence: City	Beerse	State	Country Belgium Citizenship BE
Post Office Address	c/o Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse, Belgium		
Post Office Address			
City	Turnhout	State	ZIP 2300 Country Belgium

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

Please type a plus sign (+) inside this box → ☐PTO/SB/02A (3-97)
Approved for use through 9/30/98. OMB 0651-0032
Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION**ADDITIONAL INVENTOR(S)
Supplemental Sheet**
Page 2 of 2

Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name (first and middle [if any])				Family Name or Surname			
Jasmine Elza				Viaene			
Inventor's Signature				Date			
Residence: City	Gent	State		Country	Belgium	Citizenship	BE
Post Office Address	c/o University of Gent, K.L. Ledeganckstraat 35, B-9000 Gent, Belgium						
Post Office Address							
City	Varsenare	State		ZIP	8490	Country	Belgium
Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name (first and middle [if any])				Family Name or Surname			
Marc George				Logghe			
Inventor's Signature				Date			
Residence: City	Gent	State		Country	Belgium	Citizenship	BE
Post Office Address	c/o University of Gent, K.L. Ledeganckstraat 35, B-9000 Gent, Belgium						
Post Office Address							
City	St.-Denijs-Westrem	State		ZIP	9051	Country	Belgium
Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name (first and middle [if any])				Family Name or Surname			
Jorge Eduardo <i>[Signature]</i>				Vialard <i>[Signature]</i>			
Inventor's Signature	<i>Jorge Vialard</i>			Date	03/20/01		
Residence: City	Beerse <i>[Signature]</i>	State		Country	Belgium	Citizenship	CA
Post Office Address	c/o Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse, Belgium						
Post Office Address							
City	Brussel	State		ZIP	1200	Country	Belgium

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

Please type a plus sign (+) inside this box → ☐PTO/SB/02A (3-97)
Approved for use through 9/30/98. OMB 0651-0032
Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION**ADDITIONAL INVENTOR(S)**
Supplemental Sheet
Page 1 of 2

Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor					
Given Name (first and middle [if any])				Family Name or Surname			
Bart Jozef Maria				Nelissen			
Inventor's Signature						Date	
Residence: City	Beerse	State		Country	Belgium	Citizenship	BE
Post Office Address	c/o Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse, Belgium						
Post Office Address							
City	Meerhout	State		ZIP	2450	Country	Belgium
Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor					
Given Name (first and middle [if any])				Family Name or Surname			
3-80 Marianne Denise				De Backer			
Inventor's Signature						Date	3/15/2001
Residence: City	Beerse	State		Country	Belgium	Citizenship	BE
Post Office Address	c/o Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse, Belgium						
Post Office Address							
City	San Diego	State	CA	ZIP	92122	Country	USA
Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor					
Given Name (first and middle [if any])				Family Name or Surname			
Walter Herman Maria Louis				Luyten			
Inventor's Signature						Date	
Residence: City	Beerse	State		Country	Belgium	Citizenship	BE
Post Office Address	c/o Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse, Belgium						
Post Office Address							
City	Turnhout	State		ZIP	2300	Country	Belgium

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

Please type a plus sign (+) inside this box → ☐

PTO/SB/02A (3-97)
 Approved for use through 9/30/98. OMB 0651-0032
 Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE
 Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION
ADDITIONAL INVENTOR(S)
Supplemental Sheet
 Page 2 of 2

Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor					
Given Name (first and middle (if any))		Family Name or Surname					
Jasmine Elza		Viaene					
Inventor's Signature						Date	03/22/01
Residence: City	Gent	State		Country	Belgium	Citizenship	BE
Post Office Address	c/o University of Gent, K.L. Ledeganckstraat 35, B-9000 Gent, Belgium						
Post Office Address							
City	Varsenare	State		ZIP	8490	Country	Belgium
Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor					
Given Name (first and middle (if any))		Family Name or Surname					
Marc George		Logghe					
Inventor's Signature						Date	03/22/01
Residence: City	Gent	State		Country	Belgium	Citizenship	BE
Post Office Address	c/o University of Gent, K.L. Ledeganckstraat 35, B-9000 Gent, Belgium						
Post Office Address							
City	St.-Denijs-Westrem	State		ZIP	9051	Country	Belgium
Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor					
Given Name (first and middle (if any))		Family Name or Surname					
Jorge Eduardo		Vialard					
Inventor's Signature						Date	
Residence: City	Beerse	State		Country	Belgium	Citizenship	CA
Post Office Address	c/o Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse, Belgium						
Post Office Address							
City	Brussel	State		ZIP	1200	Country	Belgium

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

DOCKET NO. JAB-1430

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants : CONTRERAS et al.

Serial No. : Art Unit:

Filed : June 4, 2001 Examiner:

For : DRUG TARGETS IN CANDIDA ALBICANS

Commissioner for Patents
Washington, D.C. 20231


ASSOCIATE POWER OF ATTORNEY

Sir:

In the matter of the above-identified application, I hereby appoint Myra H. McCormack (Reg. No. 36,602), whose postal address is One Johnson & Johnson Plaza, New Brunswick, New Jersey 08933-7003, my associate attorney to prosecute said application, to make alterations and amendments therein, to file continuing applications claiming the benefit of said application, to receive the patent and to transact all business in the Patent Office connected with said application.

I request all communications with respect to said application be addressed to Philip S. Johnson, One Johnson & Johnson Plaza, New Brunswick, New Jersey 08933-7003. All telephone calls should be directed to Myra H. McCormack (732) 524-6932.

Signed at New Brunswick, in the County of Middlesex and State of New Jersey, this 4th day of June, 2001.


Mary A. Appollina
Reg. No. 34,087
Attorney for Applicant(s)

One Johnson & Johnson Plaza
New Brunswick, NJ 08933-7003
(732) 524-3742
DATED: June 4, 2001

SEQUENCE LISTING

5 <110> Contreras, Roland
 Nelissen, Bart
 DeBacker, Marianne
 Luyten, Walter
 Viaene, Jasmine
 Logghe, Marc

10 <120> Drug Targets in Candida Albicans
 <130> 53731/000

15 <140> JAB 1430-US
 <141>
 <150> 982204122.0
 <151> 1998-12-04

20 <160> 20
 <170> PatentIn Ver. 2.0

25 <210> 1
 <211> 438
 <212> DNA
 <213> Candida albicans

30 <400> 1
 aacagctggg cttctgctaa tacattcaac ctttccata tctatactcc aacaatatga
 60
 taactgatga acaattgaat accattgcat tgacatttgg ttttgcttca ataataattaa
 120
 tcataatata tcatgccata tctactaatg tacataaatt agaagatgaa accccatcat
 180
 35 cttcatttac cagaacaaat actactgaaa ctactgttgc aagtaagaaa aagaagtaat
 240
 aactgatgga tttttcttcc taccaccaat tgaataatgc tagacttggt ggtgtgctac
 300
 40 aaatatttca aaagaaaata cgaatacttt ataaaatggg aagaacggaa gatggtttct
 360
 catttataca ctaaatacaa atcacatata catacacaaa cacaaatata tacatacacc
 420
 tatatccctt tatttgat
 45 438

<210> 2
 <211> 1380
 <212> DNA
 50 <213> Candida albicans

<400> 2
 atgttaaaaa cactaactca aactttacgc ttaactggga aagctttccc aaagggtccgt
 60
 55 ccggccttga tcagaacctc cgtgccttc gaccgttcta aacctcatgt caacattggg
 120
 actattgggc atgttgatca tggtaaaact acattgactg ctgctatcac caaagtttta
 180
 gccgaacaag gtgggtgcaa cttcttggat tatggttcta ttgatagagc tccagaagaa
 240
 60 agagctagag gtatcactat ttccactgcc cacgttgaat acgaaaccaa gaacagacac
 300

tatgcccacg ttgattgtcc aggacacgct gattatatca aaaatatgat tactgggtgcc
 360
 gctcaaattg atggtgctat cattgttgtt gctgccactg atggtcaaatt gctcaaacc
 420
 5 agagaacatt tgttattggc cagacaagtt ggtgttcaag acttggttgt gtttgtcaac
 480
 aaagtcgata ctattgatga ccctgaaatg ttggaattag tcgaaatgga aatgagagaa
 540
 ttgttatcca cctacgggtt tgatgggtgac aacactccag ttattatggg atctgcttta
 10 600
 atggcttttg aagacaagaa accagaaatt ggtaaggaag ctatcttgaa attggttagat
 660
 gctgtcgatg aacacattcc aactccatca agagacttgg aacaaccatt tttgttacca
 720
 15 gttgaagacg tgttctccat ctccggtaga ggaactgttg tcaactggtg agttgaaaga
 780
 ggtgttttga agaaggggtg agaaatcgaa attgttggtg gttttgacaa accttacaag
 840
 actactgtta ccggtattga aatgttcaaa aaagaattag actctgctat ggctgggtgac
 20 900
 aactgtggtg ttttgttaag aggtgttaaa agagatgaaa tcaagagagg tatggttttg
 960
 gccaaaccag gtactgctac ttctcacaag aagttcttgg ctcccttgta tttttgact
 1020
 25 tccgaagaag gtggtcgctt cactccattt ggtgaagggtt acaagcctca atgcttcttc
 1080
 agaactaacg atgtcactac cacattttca tcccagaag gagaagggtg tgatcattct
 1140
 caaatgatca tgccagggtg caacattgaa atggttggtg aattgatcaa atcttgtcca
 30 1200
 ttagaagtca accaacgttt caacttgaga gaagggtgta aaactgttgg tactggtttg
 1260
 attaccagaa tcatcgaata aacagaatgt gcactgtgaa taataaaaaag aaaagaggta
 1320
 35 tatataggtg actttgtatt ttgtattgaa caataaaatt ctgtaaatag taagggcctc
 1380
 <210> 3
 <211> 2283
 40 <212> DNA
 <213> Candida albicans
 <400> 3
 gaattcgccc ttaagcactc gtttcaacta tacattcagt aacaacaccc ttaatttacc
 45 60
 aaactacatt aatggaagta acacaacgga cgcagagtca gacacaacca acacaacagt
 120
 caccgacaac tcagacgcaa acccaaagca aagaggacca gaataggatt tgtcaattga
 180
 50 tttgctccac gggtcagttt ggcaattatg atttgaatat caacgataaa actatcgta
 240
 aaggtaaaat gacgtggtat tttggaagag accccaactc agatttgcaa gtggcgctcg
 300
 cgtcgagaat ttcaaacaag catTTTTcaaa tctgggtcaa cttcaatgat aaatcactat
 55 360
 ggataaagga cacttcaact aacgggacac accttaacaa cagtcgattg gtgaaaggat
 420
 caaactacct tcttaatcag ggtgatgaaa tagcagtagg ggttggtaga gacgaggacg
 480
 60 ttgtgaggtt tgctgttgc tttggtgaca aatacaaccc ggcaaagcta cctgattcga
 540

ccaacacaat taaagatgaa ggaatataca aagactttat tgtgaaaaat gaaacgatag
 600
 gccaaaggagc atttgccact gtgaaaaagg cgattgaacg atctacgggc gagtcgtacg
 660
 5 cgggtgaagat tataaatcga agaaaagcat taaataccgg tgggtggaagt gccatggcag
 720
 gagtggaccg tgaattgtcc atattagagc ggctcaacca cccaaatata gttgctctaa
 780
 10 aagcttttta tgaagatatg gacaattact atattgtgat ggaattggtg cggggcgggtg
 840
 atttgatgga ctttgtggct gcaaaccggtg caataggaga agacgcaaca caagtgatca
 900
 cgaaacagat tctagaagga attgcctatg ttcataattt aggaatctcc catcgtgatt
 960
 15 tgaagccaga taatatTTTg attatgcaag atgaccaat acttgTTaaa atcaccgact
 1020
 ttggattggc aaaattcagt gacaatctga cgtttatgaa aactTTTTgt ggtacattgg
 1080
 20 cgtatgttgc tcccgaagtt atcaccggta agtatggatc atcgcagatg gaactgcaac
 1140
 aaaaggacaa ctactcttcc ttggttgaca tttggtcttt gggatgtttg gtttatgtac
 1200
 ttttaacttc tcatttacca ttcaacggga aaaaccagca acaaatgttt gccaaagatca
 1260
 25 aaaggggcca atttcatgag gctccattaa attcatacga catttctgaa gacggaagag
 1320
 atttcttgca gtgctgcta caggTTaatc ctaaactaag gatgacggct gctgaagctt
 1380
 30 tgaacataa atggttgcaa gacttgtatg aagaggattc tgtcaaatca ttgagtttat
 1440
 cgcaatcaca gtcgcaacaa tctcgaaaga tagataatgg tatccatate gaatcattga
 1500
 gcaaaattga tgaagacggt atgcttcgtc cattggatag cgaaagaaat aggaaatcaa
 1560
 35 gtaaacagca agatttcaag gtaccaagc gtgtgattcc gttatctcaa catcctgcaa
 1620
 caccgttacc aatgtcaca cggaaaaaga ggccgtatca aatagaccct agaacaaaca
 1680
 40 aaaaagtcga tttggaagaa cctctgacaa gcaagaaagt caagctaagt gattccgttg
 1740
 ttgcggaaga ctacttgaag ttggggccac ttgcaaatto gttattccaa gaaacaataa
 1800
 atatttcaaa gtccccgttt tctttcgga gaaatgacac ttgtgattgc gagatagacg
 1860
 45 acgacagact atccaaactt catttgttca ttaccaaaga aaacgactct atatggttat
 1920
 tggataagag tactaactcg tgcttggtca acaatactag tgttgaaaa ggcaacaaag
 1980
 50 ttttgcttag aggaggggag atattacatc tcttcttga cccattgtca ctgcaacata
 2040
 taggtttcaa agtagtcctt gttgatcaac tgtctggtga acataagagt caagtggagg
 2100
 ttttgaaaca aacctcagaa gaaatgaata ttattccact tatttctggt ttaagtagta
 2160
 55 taagttcata gatttagcat atatacaagc atttctata gaaacaaagg ttcattaatt
 2220
 tagttattta cctccatgca attacattta cttcttcttc caagggcgaa ttctgcagat
 2280
 atc
 60 2283

<211> 826
<212> DNA
<213> *Candida albicans*

5 <400> 4
atgggtagta tgtgaagata caatattgaa agtggttact agaatatcta agatgtttga
60
gcccatggac atttttggat ttgataatta aaaaaagtag caatagatta ttgcgttgga
120
10 gaaagaatca ccatagttgc aagatttgat agatgttaaa atgttcacgc aggcgaaaga
180
tgtaacatct cttaaagtaa gaagaatatg gacatgaata aaaatagata gcactatatt
240
ggaacttggt gaagatatta aaatagaatg ggatttcaac atagatatc aaagtaacga
300
15 aacctcacia tcaaataaaa acaacagtaa tactaacaat tcaattttta tttttataga
360
gggtactcca tctttaggta aacgtcacia caaatctcac accttatgta acagatgtgg
420
20 ccgtcgttca ttccacgtcc aaaagaagac ctgttcttct tgtggttacc cagctgctaa
480
aatgagatct cacaactggg ccttaaaagc caaaagaaga agaactactg gtaccggtag
540
aatggcttac ttgaaacacg ttaccagaag attcaagaac gggtttccaaa ctgggtgtgc
600
25 taaagctcaa accccttccg cttaaaactaa ttactgaagt tattgggtcat gcattagtca
660
ttattcatta aagtcattgt aagcatagca aaggaagaat tggttagatt cttgttttaa
720
30 atgtaatgac tatttaatat ctgtttaaat aagagggtta gtctttattt ttttacgtat
780
acaccaaaaa aaaaagaaac aaataaaatc tgtatattaa tgttgg
826

35 <210> 5
<211> 978
<212> DNA
<213> *Candida albicans*

40 <400> 5
atgggtacta gtacaagtga agcattgaag aacatcaaaa acaaacagcg aagacagaaa
60
gtttttgcag aaataaaaca tgaaaagaat aaacaacgct ataagcaaag agccgaaaga
120
45 gctaaggaag aaagagaaaa ccccgaaata agagaggaaa gaatagcagc taatatccca
180
gatactatag atagcaaacg tatattatgat gagactatag ctgctgaagt tgaaggagat
240
gacgagtttc agtcataatt cactaacttg ttggaagaac caaagatttt gttgacaaca
300
50 agtgccaatg ctaaaaaacc ggccatgaa tttgcagaca tgatcatgga ctttttaccg
360
aatgtgacat ttatcaaaag gaagaaggaa tatacaatgc aagatatggc caaatattgc
420
55 tcgaatagag acttcaactgc attgcttgct atcaacgaag acaagaagaa ggtcaatggt
480
ataacgctca tcaatttacc tgaagggcca acattttatt ttctgattac atcaatagtt
540
gatgggaaaa gaattaaggg acacgggaaa gctgggtgatt atttacctga gattgtattg
600
60 aataatttca attcaagatt gggtaaaact gtgggaagac tatttcaaag tattttccct
660

cataaacctg aacttcaagg aagacaagtg attactttgc acaatcaacg tgattatatt
 720
 tttttcagaa gacatagata ttttttcaga aatgaggaaa aggttggatt gcaggaattg
 780
 5 ggtccgcagt ttacattaaa gctaagaaga atgcaaaagg gagtacgtgg tgatgttgtt
 840
 tgggaacaca gaccagatat ggaaagagat aagaagaagt tttatttata agcgggtgta
 900
 10 taaaggtagt agtagtgcgt ttataagtat gtgtgtgtgt ttatgcatag atgtgtaaag
 960
 agtaatacag ctaattcg
 978

 <210> 6
 15 <211> 619
 <212> DNA
 <213> Candida albicans

 <400> 6
 20 aactaatttg tttaaacatc aataccaaga agttttttaca attcaatccc acatacacca
 60
 ttaattatga attctgaaaa gattattgaa gttatcattg ctattttctt accaccagta
 120
 gctgtgttta tgaaatgtgg tgccactacc ccattatgga ttaacttggt attatgtatc
 25 180
 tttatttggt tccctgctat cttacatgcc ttatacgttg tgttgaaaga ttaaacaaac
 240
 accagagatt tactgcttga tgaattgatt actccaaaga gttgtgacta gttcccagtg
 300
 30 tgtttttttt gccttccaac tttcttttac atttttccat tactaccact gtcttcccc
 360
 ctattttgca gagttttcaa aatttatcca aaacatgtta gtcattaaac catattatta
 420
 taattattct tttttgtatt tttttccctt aaaacacggt aatttattaa tcgtttcggt
 35 480
 gtttgggtatt ttattttttt gtatttatca attggaatat atatctatac atgaatttat
 540
 tatccattgt accaattggt aaaacatttt gttagttttt tgttactagt ataaaannat
 600
 40 aataaaaagtt tanttcaac
 619

 <210> 7
 <211> 2319
 45 <212> DNA
 <213> Candida albicans

 <400> 7
 50 atgacattag gggtcgataa attcataagc aaggtcagca ctcatagacg tcaatctgaa
 60
 ccatcaatct tggaaatcgc agccaccaat tctcaaaata aatcgagaag gctaagtatg
 120
 gataatggtc attgttatgt tcgtgaatca actaataatc atcatcattt aaataccgct
 180
 55 gttgataatt tacgacagcg tgcgggatcg ttttcattta tttcacatca ccataatcac
 240
 catcagaata gtcacgataa ttatactgtc gatcccccta catcaaacgg agcacgaatt
 300
 tcccgatcac gttcacgttc caaatcagtt gggcacggag aagcaatatc accagcgtat
 360
 60 ttttccaaga ataaaaccaa agatttagtg aaacaggaaa cagcacatat cattctgaag
 420

aaattactca acatggttaca agatttggat ttacaaaacc ctattgcatt gaaaacaata
 480
 tcacaagggt cagaatcaaa gttttgtaaa atctacgtgt ctaacactaa taattgtatt
 540
 5 tacttaccag cagcaagttc aacaagtttc acttatgaag atgatgaaaa tggcggcggt
 600
 ataattgctg aagatagaaa tgatgaaatg ccaacagcag ttaataacaa tactttgtca
 660
 atggatagta taaatcattc agagactgat ttcttggtt ctcaccacc tccagattta
 720
 10 ttttctaaaa tgaaatcatt ccattcacca aattacttga cttcaaaaat cgattctgaa
 780
 tgtccaattc cacatacatt tgctgtgatt gttgaattaa ccaaggactc tttgattatt
 840
 15 aaagatcttc atttccaatt tcagtcatta actaccattt tatggccaac tggggatgca
 900
 tataatcgga ctcatgccaa ggagaaatth accattggga atatggaatg gcgtacatct
 960
 ttaagcgacg ccgactatta tatcaatagt tctaattcca acgatgttaa gctgaaaaac
 1020
 20 ttgggtcctg aagatcttat taatcgaact agagaataca aattaatcga tattgaagaa
 1080
 ccaacaatt catcaaacag tttactggat gatgacatgg atattaataa tattacgtcg
 1140
 25 ccattatcaa cgtcaccaac atcaagttca acttcaacaa attcaacctc caactcattg
 1200
 gggttcagatt catataaagc tgggtctttat gtatttttat taccaatctt attgccagaa
 1260
 catattcctg cttccattgt ttctattaat gggttcattgg ctcatacatt actgggtgaa
 1320
 30 tgcaataaat atactgataa gttgaatcgg aaatcaaaag tatcagcatc gtacaattta
 1380
 cctatgggtc gtactccacc aaacattgggt aattccattg ctgataagcc aatttatgtt
 1440
 35 aataggattt ggaatgatgc cgtacattat attataactt tccccgcaa atatgttact
 1500
 ttgggttggtg aacacatgat aaatgtgaaa ttactgccc tggtgaaaga tgtgggttact
 1560
 aagcgtatta aatttaattg attggagaga ataacttatg tttccaaaaa tttatcacga
 1620
 40 gaatatgatt atgatagtga agaccctat tgtattcatc cagtttctaa agaaaataaa
 1680
 gtacgtgaac gtgttggtgc gttatatgaa ttgaaaacga aggcaaaaca atcttctgggt
 1740
 45 ggacatcttg aagcttataa acaagaagtt atgaaatgtc cggaaaataa ctttttattt
 1800
 tcttggttatg aggttgaaaa tgataataat aacggcaacg gcaacggcaa cggcaacgga
 1860
 aacaagaacg ttaaacaana gaataaagat caaccaatga ttgctacacc tttagatatc
 1920
 50 aatgtttctt taccattttt aactactatg tctgatagtt taattatgac atcagccata
 1980
 gaagaagaag gttcagatct gcctcataca tcaagaagag ggtcggcagt gagtatgact
 2040
 55 gataataata ctacccaag taacaataac cctttatctc catttttggg agcagtggaa
 2100
 actaatggtg ctagtataaa tgaaattgggt gatcatacat tattccctga ttctaatttt
 2160
 cgacatatgg aaattaaaca tcgattacaa gttacattta ggattttctaa accggatctg
 2220
 60 gataataaaa tgcataatta tgaagtgggt attgataccc ccatcgtttt acttagttca
 2280

aaatgtcaag aagattctcc tcctccttat agttctgta
2319

5 <210> 8
<211> 255
<212> DNA
<213> Candida albicans

10 <400> 8
aacgttcgtg caaaaggcta tactgggtgat atccacgcag atgaagagca agtttaataca
60
actctttgtc aattaatgct gtacttggtt tcatttttatt tgctggcatt taaagaatac
120
ccatagttca gaaaataaaa ttgaaaaatt taaaaaaaaa cgcaatatca ttcatttttt
15 180
ttgttttttt gacaataata ttaatatgta gttaccaatg tttttagatt ttatatgttt
240
tgaaaaaata gtttg
255

20 <210> 9
<211> 119
<212> DNA
<213> Candida albicans

25 <400> 9
aaccttataca tcattataacc aactatcaaa atcataagac tcttnaactt ctgtttttga
60
tagttgggtat aatgatttat gtattatctt aattcattat tattagtttc ggtcacaaa
30 119

<210> 10
<211> 60
<212> PRT
35 <213> Candida albicans

<400> 10
Met Ile Thr Asp Glu Gln Leu Asn Thr Ile Ala Leu Thr Phe Gly Phe
1 5 10 15
40 Ala Ser Ile Ile Leu Ile Ile Ile Tyr His Ala Ile Ser Thr Asn Val
20 25 30
45 His Lys Leu Glu Asp Glu Thr Pro Ser Ser Ser Phe Thr Arg Thr Asn
35 40 45
Thr Thr Glu Thr Thr Val Ala Ser Lys Lys Lys Lys
50 55 60

50 <210> 11
<211> 426
<212> PRT
<213> Candida albicans

55 <400> 11
Met Leu Lys Thr Leu Thr Gln Thr Leu Arg Leu Thr Gly Lys Ala Phe
1 5 10 15
60 Pro Lys Val Arg Pro Ala Leu Ile Arg Thr Tyr Ala Ala Phe Asp Arg
20 25 30

Ser Lys Pro His Val Asn Ile Gly Thr Ile Gly His Val Asp His Gly
 35 40 45
 5 Lys Thr Thr Leu Thr Ala Ala Ile Thr Lys Val Leu Ala Glu Gln Gly
 50 55 60
 Gly Ala Asn Phe Leu Asp Tyr Gly Ser Ile Asp Arg Ala Pro Glu Glu
 65 70 75 80
 10 Arg Ala Arg Gly Ile Thr Ile Ser Thr Ala His Val Glu Tyr Glu Thr
 85 90 95
 Lys Asn Arg His Tyr Ala His Val Asp Cys Pro Gly His Ala Asp Tyr
 100 105 110
 15 Ile Lys Asn Met Ile Thr Gly Ala Ala Gln Met Asp Gly Ala Ile Ile
 115 120 125
 Val Val Ala Ala Thr Asp Gly Gln Met Pro Gln Thr Arg Glu His Leu
 20 130 135 140
 Leu Leu Ala Arg Gln Val Gly Val Gln Asp Leu Val Val Phe Val Asn
 145 150 155 160
 25 Lys Val Asp Thr Ile Asp Asp Pro Glu Met Leu Glu Leu Val Glu Met
 165 170 175
 Glu Met Arg Glu Leu Leu Ser Thr Tyr Gly Phe Asp Gly Asp Asn Thr
 180 185 190
 30 Pro Val Ile Met Gly Ser Ala Leu Met Ala Leu Glu Asp Lys Lys Pro
 195 200 205
 35 Glu Ile Gly Lys Glu Ala Ile Leu Lys Leu Leu Asp Ala Val Asp Glu
 210 215 220
 His Ile Pro Thr Pro Ser Arg Asp Leu Glu Gln Pro Phe Leu Leu Pro
 225 230 235 240
 40 Val Glu Asp Val Phe Ser Ile Ser Gly Arg Gly Thr Val Val Thr Gly
 245 250 255
 Arg Val Glu Arg Gly Val Leu Lys Lys Gly Glu Glu Ile Glu Ile Val
 260 265 270
 45 Gly Gly Phe Asp Lys Pro Tyr Lys Thr Thr Val Thr Gly Ile Glu Met
 275 280 285
 Phe Lys Lys Glu Leu Asp Ser Ala Met Ala Gly Asp Asn Cys Gly Val
 50 290 295 300
 Leu Leu Arg Gly Val Lys Arg Asp Glu Ile Lys Arg Gly Met Val Leu
 305 310 315 320
 55 Ala Lys Pro Gly Thr Ala Thr Ser His Lys Lys Phe Leu Ala Ser Leu
 325 330 335
 Tyr Ile Leu Thr Ser Glu Glu Gly Gly Arg Ser Thr Pro Phe Gly Glu
 340 345 350
 60 Gly Tyr Lys Pro Gln Cys Phe Phe Arg Thr Asn Asp Val Thr Thr Thr
 355 360 365

Phe Ser Phe Pro Glu Gly Glu Gly Val Asp His Ser Gln Met Ile Met
 370 375 380
 5 Pro Gly Asp Asn Ile Glu Met Val Gly Glu Leu Ile Lys Ser Cys Pro
 385 390 395 400
 Leu Glu Val Asn Gln Arg Phe Asn Leu Arg Glu Gly Gly Lys Thr Val
 405 410 415
 10 Gly Thr Gly Leu Ile Thr Arg Ile Ile Glu
 420 425
 15 <210> 12
 <211> 699
 <212> PRT
 <213> Candida albicans
 20 <400> 12
 Met Glu Val Thr Gln Arg Thr Gln Ser Gln Thr Gln Pro Thr Gln Gln
 1 5 10 15
 25 Ser Pro Thr Thr Gln Thr Gln Thr Gln Ser Lys Glu Asp Gln Asn Arg
 20 25 30
 Ile Cys Gln Leu Ile Cys Ser Thr Gly Gln Phe Gly Asn Tyr Asp Leu
 35 40 45
 30 Asn Ile Asn Asp Lys Thr Ile Val Gln Gly Lys Met Thr Trp Tyr Phe
 50 55 60
 Gly Arg Asp Pro Asn Ser Asp Leu Gln Val Ala Ser Ser Ser Arg Ile
 65 70 75 80
 35 Ser Asn Lys His Phe Gln Ile Trp Leu Asn Phe Asn Asp Lys Ser Leu
 85 90 95
 40 Trp Ile Lys Asp Thr Ser Thr Asn Gly Thr His Leu Asn Asn Ser Arg
 100 105 110
 Leu Val Lys Gly Ser Asn Tyr Leu Leu Asn Gln Gly Asp Glu Ile Ala
 115 120 125
 45 Val Gly Val Gly Arg Asp Glu Asp Val Val Arg Phe Val Val Val Phe
 130 135 140
 Gly Asp Lys Tyr Asn Pro Ala Lys Leu Pro Asp Ser Thr Asn Thr Ile
 145 150 155 160
 50 Lys Asp Glu Gly Ile Tyr Lys Asp Phe Ile Val Lys Asn Glu Thr Ile
 165 170 175
 Gly Gln Gly Ala Phe Ala Thr Val Lys Lys Ala Ile Glu Arg Ser Thr
 180 185 190
 55 Gly Glu Ser Tyr Ala Val Lys Ile Ile Asn Arg Arg Lys Ala Leu Asn
 195 200 205
 60 Thr Gly Gly Gly Ser Ala Met Ala Gly Val Asp Arg Glu Leu Ser Ile
 210 215 220

	Leu	Glu	Arg	Leu	Asn	His	Pro	Asn	Ile	Val	Ala	Leu	Lys	Ala	Phe	Tyr	
	225					230					235					240	
5	Glu	Asp	Met	Asp	Asn	Tyr	Tyr	Ile	Val	Met	Glu	Leu	Val	Pro	Gly	Gly	
					245					250					255		
	Asp	Leu	Met	Asp	Phe	Val	Ala	Ala	Asn	Gly	Ala	Ile	Gly	Glu	Asp	Ala	
				260					265					270			
10	Thr	Gln	Val	Ile	Thr	Lys	Gln	Ile	Leu	Glu	Gly	Ile	Ala	Tyr	Val	His	
			275					280					285				
	Asn	Leu	Gly	Ile	Ser	His	Arg	Asp	Leu	Lys	Pro	Asp	Asn	Ile	Leu	Ile	
15		290					295				300						
	Met	Gln	Asp	Asp	Pro	Ile	Leu	Val	Lys	Ile	Thr	Asp	Phe	Gly	Leu	Ala	
	305					310					315				320		
20	Lys	Phe	Ser	Asp	Asn	Ser	Thr	Phe	Met	Lys	Thr	Phe	Cys	Gly	Thr	Leu	
					325					330					335		
	Ala	Tyr	Val	Ala	Pro	Glu	Val	Ile	Thr	Gly	Lys	Tyr	Gly	Ser	Ser	Gln	
				340					345					350			
25	Met	Glu	Ser	Gln	Gln	Lys	Asp	Asn	Tyr	Ser	Ser	Leu	Val	Asp	Ile	Trp	
			355					360					365				
	Ser	Leu	Gly	Cys	Leu	Val	Tyr	Val	Leu	Leu	Thr	Ser	His	Leu	Pro	Phe	
30		370					375					380					
	Asn	Gly	Lys	Asn	Gln	Gln	Gln	Met	Phe	Ala	Lys	Ile	Lys	Arg	Gly	Glu	
	385				390						395					400	
35	Phe	His	Glu	Ala	Pro	Leu	Asn	Ser	Tyr	Asp	Ile	Ser	Glu	Asp	Gly	Arg	
					405					410					415		
	Asp	Phe	Leu	Gln	Cys	Cys	Leu	Gln	Val	Asn	Pro	Lys	Leu	Arg	Met	Thr	
				420					425					430			
40	Ala	Ala	Glu	Ala	Leu	Lys	His	Lys	Trp	Leu	Gln	Asp	Leu	Tyr	Glu	Glu	
			435					440					445				
	Asp	Ser	Val	Lys	Ser	Leu	Ser	Leu	Ser	Gln	Ser	Gln	Ser	Gln	Gln	Ser	
45		450					455					460					
	Arg	Lys	Ile	Asp	Asn	Gly	Ile	His	Ile	Glu	Ser	Leu	Ser	Lys	Ile	Asp	
	465				470					475					480		
50	Glu	Asp	Val	Met	Leu	Arg	Pro	Leu	Asp	Ser	Glu	Arg	Asn	Arg	Lys	Ser	
				485					490						495		
	Ser	Lys	Gln	Gln	Asp	Phe	Lys	Val	Pro	Lys	Arg	Val	Ile	Pro	Leu	Ser	
			500						505					510			
55	Gln	His	Pro	Ala	Thr	Pro	Leu	Pro	Met	Ser	Gln	Pro	Lys	Lys	Arg	Pro	
			515					520					525				
	Tyr	Gln	Ile	Asp	Pro	Arg	Thr	Asn	Lys	Lys	Val	Asp	Leu	Glu	Glu	Pro	
60		530					535					540					
	Ser	Thr	Ser	Lys	Lys	Val	Lys	Leu	Ser	Asp	Ser	Val	Val	Ala	Glu	Asp	
	545					550				555						560	

	Tyr	Leu	Lys	Leu	Gly	Pro	Leu	Ala	Asn	Ser	Leu	Phe	Gln	Glu	Thr	Ile	
					565					570						575	
5	Asn	Ile	Ser	Lys	Ser	Pro	Phe	Ser	Phe	Gly	Arg	Asn	Asp	Thr	Cys	Asp	
				580					585					590			
	Cys	Glu	Ile	Asp	Asp	Asp	Arg	Leu	Ser	Lys	Leu	His	Cys	Val	Ile	Thr	
			595					600					605				
10	Lys	Glu	Asn	Asp	Ser	Ile	Trp	Leu	Leu	Asp	Lys	Ser	Thr	Asn	Ser	Cys	
		610					615					620					
15	Leu	Val	Asn	Asn	Thr	Ser	Val	Gly	Lys	Gly	Asn	Lys	Val	Leu	Leu	Arg	
	625					630					635					640	
	Gly	Gly	Glu	Ile	Leu	His	Leu	Phe	Phe	Asp	Pro	Leu	Ser	Ser	Gln	His	
					645					650					655		
20	Ile	Gly	Phe	Lys	Val	Val	Leu	Val	Asp	Gln	Ser	Ser	Gly	Glu	His	Lys	
				660					665					670			
	Ser	Gln	Val	Glu	Val	Leu	Lys	Gln	Thr	Ser	Glu	Glu	Met	Asn	Ile	Ile	
			675					680					685				
25	Pro	Leu	Ile	Ser	Gly	Leu	Ser	Ser	Ile	Ser	Ser						
		690					695										
30	<210> 13 <211> 295 <212> PRT <213> Candida albicans																
35	<400> 13 Met Gly Thr Ser Thr Ser Glu Ala Leu Lys Asn Ile Lys Asn Lys Gln 1 5 10 15																
40	Arg	Arg	Gln	Lys	Val	Phe	Ala	Glu	Ile	Lys	His	Glu	Lys	Asn	Lys	Gln	
				20					25					30			
	Arg	His	Lys	Gln	Arg	Ala	Glu	Arg	Ala	Lys	Glu	Glu	Arg	Glu	Asn	Pro	
			35					40					45				
45	Glu	Leu	Arg	Glu	Glu	Arg	Ile	Ala	Ala	Asn	Ile	Pro	Asp	Thr	Ile	Asp	
		50					55					60					
	Ser	Lys	Arg	Ile	Tyr	Asp	Glu	Thr	Ile	Ala	Ala	Glu	Val	Glu	Gly	Asp	
	65					70					75					80	
50	Asp	Glu	Phe	Gln	Ser	Tyr	Phe	Thr	Asn	Leu	Leu	Glu	Glu	Pro	Lys	Ile	
					85					90					95		
	Leu	Leu	Thr	Thr	Ser	Ala	Asn	Ala	Lys	Lys	Pro	Ala	Tyr	Glu	Phe	Ala	
55				100					105					110			
	Asp	Met	Ile	Met	Asp	Phe	Leu	Pro	Asn	Val	Thr	Phe	Ile	Lys	Arg	Lys	
			115					120					125				
60	Lys	Glu	Tyr	Thr	Met	Gln	Asp	Met	Ala	Lys	Tyr	Cys	Ser	Asn	Arg	Asp	
		130					135					140					

Phe Thr Ala Leu Leu Val Ile Asn Glu Asp Lys Lys Lys Val Asn Gly
 145 150 155 160
 5 Ile Thr Leu Ile Asn Leu Pro Glu Gly Pro Thr Phe Tyr Phe Ser Ile
 165 170 175
 Thr Ser Ile Val Asp Gly Lys Arg Ile Lys Gly His Gly Lys Ala Gly
 180 185 190
 10 Asp Tyr Leu Pro Glu Ile Val Leu Asn Asn Phe Asn Ser Arg Leu Gly
 195 200 205
 Lys Thr Val Gly Arg Leu Phe Gln Ser Ile Phe Pro His Lys Pro Glu
 210 215 220
 15 Leu Gln Gly Arg Gln Val Ile Thr Leu His Asn Gln Arg Asp Tyr Ile
 225 230 235 240
 Phe Phe Arg Arg His Arg Tyr Ile Phe Arg Asn Glu Glu Lys Val Gly
 245 250 255
 20 Leu Gln Glu Gly Pro Gln Phe Thr Leu Lys Leu Arg Arg Met Gln Lys
 260 265 270
 25 Gly Val Arg Gly Asp Val Val Trp Glu His Arg Pro Asp Met Glu Arg
 275 280 285
 Asp Lys Lys Lys Phe Tyr Leu
 290 295
 30
 <210> 14
 <211> 55
 <212> PRT
 35 <213> Candida albicans
 <400> 14
 Met Asn Ser Glu Lys Ile Ile Glu Val Ile Ile Ala Ile Phe Leu Pro
 1 5 10 15
 40 Pro Val Ala Val Phe Met Lys Cys Gly Ala Thr Thr Pro Leu Trp Ile
 20 25 30
 45 Asn Leu Val Leu Cys Ile Phe Ile Trp Phe Pro Ala Ile Leu His Ala
 35 40 45
 Leu Tyr Val Val Leu Lys Asp
 50 55
 50
 <210> 15
 <211> 773
 <212> PRT
 <213> Candida albicans
 55
 <400> 15
 Met Thr Leu Gly Phe Asp Lys Phe Ile Ser Lys Val Ser Thr His Arg
 1 5 10 15
 60 Arg Gln Ser Glu Pro Ser Ile Leu Glu Ile Ala Ala Thr Asn Ser Gln
 20 25 30

Asn Lys Ser Arg Arg Leu Ser Met Asp Asn Gly His Cys Tyr Val Arg
 35 40 45
 5 Glu Ser Thr Asn Asn His His His Leu Asn Thr Val Val Asp Asn Leu
 50 55 60
 Arg Gln Arg Ala Gly Ser Phe Ser Phe Ile Ser His His His Asn His
 65 70 75 80
 10 His Gln Asn Ser His Asp Asn Tyr Thr Val Asp Pro Leu Thr Ser Asn
 85 90 95
 Gly Ala Arg Ile Ser Arg Ser Arg Ser Arg Ser Lys Ser Val Gly His
 100 105 110
 15 Gly Glu Ala Ile Ser Pro Ala Tyr Phe Ser Lys Asn Lys Thr Lys Asp
 115 120 125
 Leu Val Lys Gln Glu Thr Ala His Ile Ile Ser Lys Lys Leu Leu Asn
 130 135 140
 Met Leu Gln Asp Leu Asp Leu Gln Asn Pro Ile Ala Leu Lys Thr Ile
 145 150 155 160
 25 Ser Gln Gly Ser Glu Ser Lys Phe Cys Lys Ile Tyr Val Ser Asn Thr
 165 170 175
 Asn Asn Cys Ile Tyr Leu Pro Ala Ala Ser Ser Thr Ser Phe Thr Tyr
 180 185 190
 30 Glu Asp Asp Glu Asn Gly Gly Val Ile Ile Ala Glu Asp Arg Asn Asp
 195 200 205
 35 Glu Met Pro Thr Ala Val Asn Asn Asn Thr Leu Ser Met Asp Ser Ile
 210 215 220
 Asn His Ser Glu Thr Asp Phe Ser Asp Ser Pro Pro Pro Pro Asp Leu
 225 230 235 240
 40 Phe Ser Lys Met Lys Ser Phe His Ser Pro Asn Tyr Leu Thr Ser Lys
 245 250 255
 Ile Asp Ser Glu Cys Pro Ile Pro His Thr Phe Ala Val Ile Val Glu
 260 265 270
 45 Leu Thr Lys Asp Ser Leu Ile Ile Lys Asp Leu His Phe Gln Phe Gln
 275 280 285
 Ser Leu Thr Thr Ile Leu Trp Pro Thr Gly Asp Ala Tyr Asn Arg Thr
 290 295 300
 His Ala Lys Glu Lys Phe Thr Ile Gly Asn Met Glu Trp Arg Thr Ser
 305 310 315 320
 55 Leu Ser Asp Ala Asp Tyr Tyr Ile Asn Ser Ser Asn Ser Asn Asp Val
 325 330 335
 Lys Ser Lys Asn Leu Gly Pro Glu Asp Leu Ile Asn Arg Thr Arg Glu
 340 345 350
 60 Tyr Lys Leu Ile Asp Ile Glu Glu Pro Asn Asn Ser Ser Asn Ser Leu
 355 360 365

Ser Asp Asp Asp Met Asp Ile Asn Asn Ile Thr Ser Pro Leu Ser Thr
 370 375 380
 5 Ser Pro Thr Ser Ser Ser Thr Ser Thr Asn Ser Thr Ser Asn Ser Leu
 385 390 395 400
 Gly Ser Asp Ser Tyr Lys Ala Gly Leu Tyr Val Phe Leu Leu Pro Ile
 405 410 415
 10 Leu Leu Pro Glu His Ile Pro Ala Ser Ile Val Ser Ile Asn Gly Ser
 420 425 430
 Leu Ala His Thr Leu Ser Val Glu Cys Asn Lys Tyr Thr Asp Lys Leu
 435 440 445
 15 Asn Arg Lys Ser Lys Val Ser Ala Ser Tyr Asn Leu Pro Met Val Arg
 450 455 460
 20 Thr Pro Pro Asn Ile Gly Asn Ser Ile Ala Asp Lys Pro Ile Tyr Val
 465 470 475 480
 Asn Arg Ile Trp Asn Asp Ala Val His Tyr Ile Ile Thr Phe Pro Arg
 485 490 495
 25 Lys Tyr Val Thr Leu Gly Cys Glu His Met Ile Asn Val Lys Leu Ser
 500 505 510
 30 Pro Met Val Lys Asp Val Val Ile Lys Arg Ile Lys Phe Asn Val Leu
 515 520 525
 Glu Arg Ile Thr Tyr Val Ser Lys Asn Leu Ser Arg Glu Tyr Asp Tyr
 530 535 540
 35 Asp Ser Glu Asp Pro Tyr Cys Ile His Pro Val Ser Lys Glu Asn Lys
 545 550 555 560
 Val Arg Glu Arg Val Val Ser Leu Tyr Glu Leu Lys Thr Lys Ala Lys
 565 570 575
 40 Gln Ser Ser Gly Gly His Leu Glu Ala Tyr Lys Gln Glu Val Met Lys
 580 585 590
 Cys Pro Glu Asn Asn Leu Leu Phe Ser Cys Tyr Glu Val Glu Asn Asp
 595 600 605
 45 Asn Asn Asn Gly Asn Gly Asn Gly Asn Gly Asn Gly Asn Lys Asn Val
 610 615 620
 50 Lys Gln Lys Asn Lys Asp Gln Pro Met Ile Ala Thr Pro Leu Asp Ile
 625 630 635 640
 Asn Val Ser Leu Pro Phe Leu Thr Thr Met Ser Asp Ser Leu Ile Met
 645 650 655
 55 Thr Ser Ala Ile Glu Glu Glu Gly Ser Asp Ser Pro His Thr Ser Arg
 660 665 670
 60 Arg Gly Ser Ala Val Ser Met Thr Asp Asn Asn Thr Thr Pro Ser Asn
 675 680 685
 Asn Asn Pro Leu Ser Pro Phe Leu Gly Ala Val Glu Thr Asn Gly Ala

690 695 700
 Ser Ile Asn Glu Ile Gly Asp His Thr Leu Phe Pro Asp Ser Asn Phe
 705 710 715 720
 5 Arg His Ile Glu Ile Lys His Arg Leu Gln Val Thr Phe Arg Ile Ser
 725 730 735
 10 Lys Pro Asp Ser Asp Asn Lys Met His His Tyr Glu Val Val Ile Asp
 740 745 750
 Thr Pro Ile Val Leu Leu Ser Ser Lys Cys Gln Glu Asp Ser Pro Pro
 755 760 765
 15 Pro Tyr Ser Ser Val
 770
 20 <210> 16
 <211> 90
 <212> PRT
 <213> Candida albicans
 25 <400> 16
 Met Gly Glu Gly Thr Pro Ser Leu Gly Lys Arg His Asn Lys Ser His
 1 5 10 15
 Thr Leu Cys Asn Arg Cys Gly Arg Arg Ser Phe His Val Gln Lys Lys
 20 25 30
 30 Thr Cys Ser Ser Cys Gly Tyr Pro Ala Ala Lys Met Arg Ser His Asn
 35 40 45
 35 Trp Ala Leu Lys Ala Lys Arg Arg Arg Thr Thr Gly Thr Gly Arg Met
 50 55 60
 Ala Tyr Leu Lys His Val Thr Arg Arg Phe Lys Asn Gly Phe Gln Thr
 65 70 75 80
 40 Gly Val Ala Lys Ala Gln Thr Pro Ser Ala
 85 90
 45 <210> 17
 <211> 19
 <212> DNA
 <213> artificial sequence
 50 <220>
 <223> Description of Artificial Sequence:primer
 <400> 17
 tgcagctcga cctcgactg 19
 55 <210> 18
 <211> 19
 <212> DNA
 <213> artificial sequence
 60 <220>
 <223> Description of Artificial Sequence:primer

```

<400> 18
gcgtgaatgt aagcgtgac 19

5
<210> 19
<211> 20
<212> DNA
<213> artificial sequence

10
<220>
<223> Description of Artificial Sequence:primer

<400> 19
15 tgagcagctc gccgtcgcgc 20

<210> 20
<211> 22
20 <212> DNA
<213> artificial sequence

<220>
<223> Description of Artificial Sequence:primer

25
<400> 20
gagttatacc ctgcagctcg ac 22

30
<210> 21
<211> 6671
<212> DNA
<213> Artificial Sequence

35
<220>
<223> Description of Artificial Sequence:DNA plasmid

<400> 21
40 ttccatcggg gaaagtgggg gggaaaaaat tttaagcagt tcacaaaacc ttccaaaaaa
60
tatatggaca aagatgattg tattttcccg acacaaaaat cataattaat tatgagaaag
120
ttaaatgtaa cgttacaatt tatgtttatt tgaaggtgaa aagcgattta tgatttttcc
180
45 gaaatgaaaa ttttttttag gttttttttt tttgtcgggc aaagaaaaac tgaacaagga
240
ttattaaaaat ttttggtggt tgtttgtgtc tggagaattc attcctctct catcttcaca
300
caatgttttag acatctgaca cgattcatga tagttcgggt tccgggggtg gtgttttagtt
50 360
ttcgtttttc tttttttttg gaaagaatgt tttagctcat tggttttott tcttcattca
420
atagttttga aagaatttgc ccacttggtt ttacaatcat ataaaattaa actttgatat
480
55 aaaatagagt ttgaaagttt cccagatcct ttttgatttc tttgtaaatt tttttttctc
540
ccacatatac acacatacaa accgattttt ataagaaaga gttataccct gcagctcgac
600
ctcgactggt taaacctgca ggcattgcaag cttggccaaa aaggcctcga ggaacatgac
660
60 caacaagtgt ctctccaaa ttgctctcct gttgtgcttc tccactacag ctctttccat
720

```

gagctacaac ttgcttgat tectacaaag aagcagcaat tttcagtgtc agaagctcct
 780
 gtggcaattg aatgggaggc ttgaatactg cctcaaggac aggatgaact ttgacatccc
 840
 5 tgaggagatt aagcagctgc agcagttcca gaaggaggac gccgcattga ccatctatga
 900
 gatgctccag aacatctttg ctattttcag acaagattca tctagcactg gctggaatga
 960
 10 gactattggt gagaacctcc tggctaattg ctatcatcag ataaaccatc tgaagacagt
 1020
 cctggaagaa aaactggaga aagaagattt caccagggga aaactcatga gcagtctgca
 1080
 cctgaaaaga tattatggga ggattctgca ttacctgaag gccaaggagt acagtcactg
 1140
 15 tgcctggacc atagtcagag tggaaatcct aaggaaacttt tacttcatta acagacttac
 1200
 aggttacctc cgaaactgaa gatctcctag cctgtgcctc tgggactgga caattgcttc
 1260
 aagcattctt caaccagcag atgctgttta agtgactgat ggctaattgta ctgcatatga
 1320
 20 aaggacacta gaagattttg aaatttttat taaattatga gttattttta tttattttaa
 1380
 tttatttttg gaaaataaat ttttttgggt gcaaaagtcc ctcgaggcct agcgggcgcc
 1440
 25 tagaggatcc ccgggcgcta ggccggcgct aggccttttt ggccgaattc gagctcggtg
 1500
 cccggggaga tccgtcccc ttttcctttg tcgatatcat gtaattagtt atgtcacgct
 1560
 30 tacattcacg cctccccccc acatccgctc taaccgaaaa ggaaggagtt agacaacctg
 1620
 aagtctaggt ccttatttat ttttttatag ttatgttagt attaagaacg ttattttat
 1680
 ttcaaatttt tctttttttt ctgtacagac gcgtgtacgc atgtaacatt atactgaaaa
 1740
 35 ccttgcttga gaaggttttg ggacgctcga aggctttaat ttgcaagcta gcttggcgta
 1800
 atcatggta tagctgtttc ctgtgtgaaa ttgttatccg ctacacaattc cacacaacat
 1860
 acgagccgga agcataaagt gtaaagcctg gggcgcttaa tgagtgaagt aactcacatt
 1920
 40 aattgcgttg cgctcactgc ccgctttcca gtcgggaaac ctgtcgtgcc agagatctct
 1980
 gcattaatga atcggccaac gcgcggggag aggcggtttg cgtattgggc gctcttcgac
 2040
 45 ttctcgtc actgactcgc tgcgctcggg cgttcggctg cggcgagcgg tatcagatcg
 2100
 atctcactca aaggcggtta tacggttatc cacagaatca ggggataacg caggaaagaa
 2160
 catgtgagca aaaggccagc aaaaggccag gaaccgtaaa aaggccgcgt tgctggcggt
 2220
 50 tttccatagg ctccgcccc ctgacgagca tcacaaaaat cgacgctcaa gtcagaggtg
 2280
 gcgaaacccg acaggactat aaagatacca ggctttccc cctggaagct cctcgtgacg
 2340
 55 ctctcctggt ccgacctgc cgtttaccgg atacctgtcc gcctttctcc cttcggaag
 2400
 cgtggcgctt tctcatagct cacgctgtag gtatctcagt tcggtgtagg tcgttcgctc
 2460
 caagctgggc tgtgtgcacg aacccccgt tcagcccgac cgctgcgctt tatccggtta
 2520
 60 ctatcgtctt gagtccaacc cggtaagaca cgacttatcg ccaactggcag cagccactgg
 2580

taacaggatt agcagagcga ggtatgtagg cggtgctaca gagttcttga agtgggtggcc
2640
taactacggc tacactagaa ggacagtatt tggatatctgc gctctgctga agccagttac
2700
5 cttcggaaaa agagttggta gctcttgatc cggcaaacaa accaccgctg gtagcgggtg
2760
tttttttggt tgcaagcagc agattacgcg cagaaaaaaaa ggatctcaag aagatccttt
2820
gatcttttct acgggggtctg acgctcagtg gaacgaaaaac tcacgttaag ggatttttgg
2880
10 catgagatta tcaaaaagga tcttcaccta gatcctttta aattaaaaat gaagttttta
2940
atcaatctaa agaagtggat ctaggaaaat ttaattttta cttcaaaatt tagttagatt
3000
15 agtatatatg agtaaacttg gtctgacagt taccaatgct taatcagtga ggcacctatc
3060
tcagcgatct gtctatttgc ttcattccata gttgcctgac tccccgctgt gtagataact
3120
acgatacggg agggccttacc atctggcccc agtgctgcaa tgataccgcg agaccacgc
3180
20 tcaccggctc cagattttatc agcaataaac cagccagccg gaagggccga gcgcagaagt
3240
ggtcctgcaa cttttatccgc ctccatccag tctattaatt gttgccggga agctagagta
3300
25 agtagttcgc cagttaatag tttgcgcaac gttgttgcca ttgctacagg catcgtgggtg
3360
tcacgctcgt cgtttgggat ggcttcattc agctccggtt cccaacgata aaggcgagtt
3420
acatgatccc ccatgttgtg caaaaaagcg gttagctcct tcggctcctc gatcgttgtc
3480
30 agaagtaagt tggccgcagt gttatcactc atggttatgg cagcactgca taattctctt
3540
actgtcatgc catccgtaag atgcttttct gtgactgggtg agtactcaac caagtcattc
3600
35 tgagaatagt gtatgcggcg accgagttgc tcttgcccgg cgtcaatacg ggataatacc
3660
gcgccacata gcagaacttt aaaagtgtc atcattggaa aacgttcttc ggggcgaaaa
3720
ctctcaagga tcttaccgct gttgagatcc agttcgatgt aaccactcgc tgcacccaac
3780
40 tgatcttcag catcttttac tttcaccagc gtttctgggt gagcaaaaac aggaaggcaa
3840
aatgccgcaa aaaaggaat aagggcgaca cggaaatgtt gaatactcat actcttcctt
3900
45 tttcaatatt attgaagcat ttatcagggt tattgtctca tgagcggata catatttgaa
3960
tgtatttaga aaaataaaca aataggggtt ccgcgcacat tccccgaaa agtgccacct
4020
gacgtctaag aaaccattat tatcatgaca ttaacctata aaaataggcg tatcacgagg
4080
50 ccttttcgtc tcgcgcgttt cgggtgatgac ggtgaaaacc tctgacacat gcagctcccc
4140
gagacggtea cagcttgtct gtaagcggat gccgggagca gacaagcccg tcagggcgcg
4200
55 tcagcgggtg ttggcgggtg tcggggctgg cttaactatg cggcatcaga gcagattgta
4260
ctgagagtgc accatatcga cgctctccct tatgcgactc ctgcattagg aagcagccca
4320
gtagtaggtt gaggccgttg agcaccgccg ccgcaaggaa tgggtgcatgc aaggagatgg
4380
60 cgcccaacag tccccgggc acggggcctg ccaccatacc cagccgaaa caagcactaa
4440

taggaattga tttggatggt ataaacggaa acaaaaaaaaa gagctggtac tactttcttt
 4500
 aaaattatatt tattatttga ttttatttaa tagtatatat tatattttga acgtagatta
 4560
 5 ttttggtgaa agttgctgta gtgccattga ttcgtaacac taattctgta ttagtcattc
 4620
 ctcttggttg atagtatcca aaaaaacggc ttttttttg caatcttatt tcctgcatat
 4680
 10 tatacagata acataatgaa agaaaaaatc ttttttttg ttcttcaatg atgatttcaa
 4740
 ccattctttt aaacattgat caattcctga gcaacaaccc catcacact ggtttatata
 4800
 ccgccccttt tacagttgaa gaaagaaata gaaatagaaa tagcaaaca aagatatgac
 4860
 15 agtcaacact aagacctata gtgagagagc agaaactcat gcctcaccag tagcacagcg
 4920
 attatttctga ttaatggaac tgaagaaaac caatttatgt gcatcaattg acgttgatac
 4980
 20 cactaaggag ttcctcgagt taattgataa attaggtcct tatgtatgct taatcaagac
 5040
 tcatattgat ataataatg atttttctta tgaatccact attgaaccat tattagaact
 5100
 ttcacgtaaa catcaattta tgatttttga agatagaaaa tttgctgata ttggtataac
 5160
 25 cgtaaagaaa caatatattg gtggagttta taaaattagt agttgggcag atattaccaa
 5220
 tgctcatggt gtcactggga atggagtggt tgaaggatta aaacaggag ctaaagaaac
 5280
 30 caccaccaac caagagccaa gaggggtatt gatgttagct gaattatcat cagtgggac
 5340
 attagcatat ggagaatatt ctcaaaaaac tgttgaaatt gctaaatccg ataaggaatt
 5400
 tgttattgga tttattgccc aacgtgatat ggggtggcaa gaagaaggat ttgattggct
 5460
 35 tattatgaca cctggagttg gattagatga taaaggtgat ggattaggac aacaatatag
 5520
 aactgttgat gaagttgtta gcaactggaac tgatattatc attgttggtta gaggattggt
 5580
 40 tggtaaagga agagatccag atattgaagg taaaaggat agaaatgctg gttggaatgc
 5640
 ttatttgaaa aagactggcc aattataaat gtgaaggggg agattttcac tttattagat
 5700
 ttgtatatat gtagaataaa taaataaata agttaataa ataattaaat aagggtggta
 5760
 45 attattacta ttacaatca aagggtggtc ttctagctgt aatccgggca gcgcaacgga
 5820
 acattcatca gtgtaaaaat ggaatcaata aagccctgcg ctcatgagcc cgaagtggcg
 5880
 50 agcccgatct tccccatcgg tgatgtcggc gatataggcg ccagcaaccg cacctgtggc
 5940
 gccgcagcgc gcagggtcag cctgaatacg cgtttaatga ccagcacagt cgtgatggca
 6000
 aggtcagaat agcccaagtc ggccgagggg cctgtacagt gaggaagat ctgatattga
 6060
 55 cgaagaggaa ccaatgtaac gttacactga agaaaacaca caataaacgg gaagaaacgg
 6120
 tgtaaaagtg tgaaaataat ttttgaatat catttccctt ggtttaattc caaacgaaac
 6180
 gtgttttttt tagagaatgg gaattcttat tggatgtcta gattgtttgt ttactccaga
 6240
 60 ctgtgcacaa aaacgttttg atggatgatc agaagatatt ttaggctta gctctaaata
 6300

taagaaatga tgcttgaaaa accagacaga aattgagttt caaaaattgg taatgtgagg
 6360
 tattagtcaa ctaaccaa at acaatgcaa accggttgat acatttcatt ttgaaaataa
 6420
 5 tgaaactgga attggatgac cagcacacaa acacataaag taattatggg aattagaagc
 6480
 gaacatagag gagtacttgg ccacgaacag aatacaagtg ggaacactat tttctccatt
 6540
 10 gtttttagttc tgtttttttg tcagcctagt tttgtgctat gtgtaaaaaa tattgccaaag
 6600
 aaaaaaagct tgttttgtgg ccagtgtccg aaaaaaattt tggggaatct tcggattaat
 6660
 ttatgttttc a
 6671
 15
 <210> 22
 <211> 7127
 <212> DNA
 20 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence:DNA plasmid
 25 <400> 22
 agcttgagta ttctatagtg tcacctaaat agcttggcgt aatcatggtc atagctgttt
 60
 cctgtgtgaa attgttatcc gtcacaatt ccacacaaca tacgagccgg aagcataaag
 120
 30 tgtaaagcct ggggtgccta atgagtgagc taactcacat taattgcgtt gcgctcactg
 180
 cccgctttcc agtcgggaaa cctgtcgtgc cagctgcatt aatgaatcgg ccaacgcgcg
 240
 gggagaggcg gtttgcgtat tgggcgctct tccgcttccg cgctcactga ctcgctgcgc
 300
 35 tcggtcgttc ggctgcggcg agcggatatca gctcactcaa aggcggtaat acggttatcc
 360
 acagaatcag gggataacgc aggaagaac atgtgagcaa aaggccagca aaaggccagg
 420
 40 aaccgtaaaa aggccgcgtt gctggcggtt ttccataggg tccgcccccc tgacgagcat
 480
 cacaaaaatc gacgctcaag tcagaggtgg cgaaaccga caggactata aagataccag
 540
 gcgtttcccc ctggaagctc cctcgtgcgc tctcctgttc cgaccctgcc gcttaccgga
 600
 45 tacctgtccg cttttctccc ttcgggaagc gtggcgcttt ctcatagctc acgctgtagg
 660
 tatctcagtt cgggtgtaggt cgttcgctcc aagctgggct gtgtgcacga accccccggt
 720
 50 cagcccagacc gctgcgcctt atccggtaac tatcgtcttg agtccaaccc ggtaagacac
 780
 gacttategc cactggcagc agccactggg aacaggatta gcagagcgag gtatgtaggc
 840
 ggtgctacag agttcttgaa gtggtggcct aactacggct acactagaag gacagtattt
 900
 55 ggtatctgcg ctctgctgaa gccagttacc ttcggaaaaa gagttggtag ctcttgatcc
 960
 ggcaaaaaa ccaccgctgg tagcggtggt tttttgttt gcaagcagca gattacgcgc
 1020
 60 agaaaaaaag gatctcaaga agatcctttg atcttttcta cggggctctga cgctcagtgg
 1080


aacgaaaact cacgttaagg gatttttggtc atgagattat caaaaaggat cttcacctag
 1140
 atccttttaa attaaaaatg aagtttttaa tcaatctaaa gtatatatga gtaaacttgg
 1200
 5 tctgacagtt accaatgctt aatcagtgag gcacctatct cagcgatctg tctatttcgt
 1260
 tcatccatag ttgcctgact ccccgctcgtg tagataacta gaccacgct caccggctcc
 1320
 10 agatttatca gcaataaacc agccagccgg aagggcggag cgcagaagtg gtcttgcaac
 1380
 tttatccgcc tccatccagt ctattaattg ttgcgggaa gctagagtaa gtagttcgcc
 1440
 agttaatagt ttgcgcaacg ttgttgccat tgctacaggc atcgtggtgt cagctcgtc
 1500
 15 gtttgggatg gtttcattca gtcgggttc ccaacgatca aggcgagtta catgatcccc
 1560
 catgttgtgc aaaaaagcgg ttagctcctt cggctcctcg atcgttgtca gaagtaagtt
 1620
 20 ggccgcagtg ttatcactca tggttatggc agcactgcat aattctctta ctgtcatgcc
 1680
 atccgtaaga tgcttttctg tgactggtga gtactcaacc aagtcattct gagaatagtg
 1740
 tatggggcga ccgagttgct cttgcccggc gtcaatacgg gataataccg cgccacatag
 1800
 25 cagaacttta aaagtgctca tcattggaaa acgttcttcg gggcgaaaac tctcaaggat
 1860
 cttaccgctg ttgagatcca gtctgatgta acccactcgt gcacccaact gatcttcagc
 1920
 30 atcttttact ttcaccagcg tttctgggtg agcaaaaaca ggaaggcaaa atgccgcaaa
 1980
 aaaggggaata agggcgacac ggaaatgttg aatactcata ctcttccttt ttcaatatta
 2040
 ttgaagcatt tatcaggggt attgtctcat gagcggatac atatttgaat gtatttagaa
 2100
 35 aaataaacia ataggggttc cgcgacatt tccccgaaaa gtgccacctg acgtctaaga
 2160
 aaccattatt atcatgacat taacctataa aaataggcgt atcacgagc ctttctgtct
 2220
 40 cgcgcggttc ggtgatgacg gtgaaaacct ctgacacatg cagctcccgg agacggctac
 2280
 agcttgtctg taagcggatg ccgggagcag acaagcccgt cagggcgctg cagcgggtgt
 2340
 tggcgggtgt cggggctggc ttaactatgc ggcacagag cagattgtac tgagagtga
 2400
 45 ccatatgcgg tgtgaaatac cgcacagatg cgtaaggaga aaataccgca tcaggcgaaa
 2460
 ttgtaaactg taatattttg ttaaaattcg cgtaaataat ttgttaaata agctcatttt
 2520
 50 ttaaccaata ggccgaaatc ggcaaaatcc cttataaatc aaaagaatag accgagatag
 2580
 ggttgagtgt tgttccagtt tggaacaaga gtccactatt aaagaacgtg gactccaacg
 2640
 tcaaagggcg aaaaaccgtc tatcagggcg atggcccact acgtgaacca tcacccaat
 2700
 55 caagtttttt gcggtcgagg tgccgtaaag ctctaaatcg gaaccctaaa gggagcccc
 2760
 gatttagagc ttgacgggga aagccggcga acgtggcgag aaaggaaggg aagaaagcga
 2820
 aaggagcggg cgctagggcg ctggcaagtg tagcggtcac gctgcgcgta accaccacac
 2880
 60 ccgcccgcgt taatgcgccg ctacagggcg cgtccattcg ccattcaggc tgcgcaactg
 2940

ttgggaaggg cgatcgggtgc gggcctcttc gctattacgc cagctggcga aagggggatg
 3000
 tgctgcaagg cgattaagtt gggtaacgcc aggggttttcc cagtcacgac gttgtaaaac
 3060
 5 gacggccagt gaattgtaat acgactcaact atagggcgaa ttgggttttcc aatgatgagc
 3120
 acttttaaag ttctgctatg tggcgcggtg ttatcccgtg ttgacgccgg gcaagagcaa
 3180
 10 ctcggtcgcc gcatacacta ttctcagaat gacttggttg agtactaata ggaattgatt
 3240
 tggatggtat aaacggaaac aaaaaaaga gctgggtacta ctttctttaa aattatttta
 3300
 ttatttgatt ttatttaata gtatatatta tattttgaac gtagattatt ttgttgaaag
 3360
 15 ttgctgtagt gccattgatt cgtaacacta attctgtatt agtcattcct cttgtttgat
 3420
 agtatccaaa aaaacggcta tttttttgca atcttatttc ctgcatatta tacagataac
 3480
 20 ataataaag aaaaaatctt tttttttggt cttcaatgat gatttcaacc attcttttaa
 3540
 acattgatca attcctgagc aacaacccca tacacactgg tttatatacc gccctttta
 3600
 cagttgaaga aagaaataga aatagaaata gcaaacaaaa gatatgacag tcaacactaa
 3660
 25 gacctatagt gagagagcag aaactcatgc ctccaccagta gcacagcgat tatttcgatt
 3720
 aatggaactg aagaaaacca atttatgtgc atcaattgac gttgatacca ctaaggaatt
 3780
 30 ccttgaatta attgataaat taggtcctta tgtatgctta atcaagactc atattgatat
 3840
 aatcaatgat ttttcctatg aatccactat tgaaccatta ttagaacttt cacgtaaaca
 3900
 tcaatttatg atttttgaag atagaaaatt tgctgatatt ggtaataaccg taaagaaaca
 3960
 35 atatattggt ggagtttata aaattagtag ttgggcagat attaccaatg ctcatggtgt
 4020
 cactgggaat ggagtgggtg aaggattaaa acagggagct aaagaaacca ccaccaacca
 4080
 40 agagccaaga gggttattga tgtagctga attatcatca gtgggatcat tagcatatgg
 4140
 agaataattct caaaaaactg ttgaaattgc taaatccgat aaggaatttg ttattggatt
 4200
 tattgcccga cgtgatatgg gtggccaaga agaaggattt gattggctta ttatgacacc
 4260
 45 tggagttaga ttagatgata aaggatgatg attaggacaa caatatagaa ctgttgatga
 4320
 agttgttagc actggaactg atattatcat tgttggtaga ggattgtttg gtaaaggaag
 4380
 50 agatccagat attgaaggta aaaggatatag aaatgctggt tggaatgctt atttgaaaaa
 4440
 gactggccaa ttataaatgt gaagggggag attttcactt tattagattt gtatatatgt
 4500
 agaataaata aataaataag ttaaataaat aattaaataa ggggtggaat tattactatt
 4560
 55 tacaatcaaa ggtggtcctt ctagctgtaa tccgggcagc gcaacggaac attcatcagt
 4620
 gtaaaaaatgg aatcaataaa gccctgcgca gcgcgcaggg tcagcctgaa tacgcgttta
 4680
 60 atgaccagca cagtcgtgat ggcaaggtca gaatagccca agtcggccga ggggcctgta
 4740
 cagtgagga agatctgata ttgacgaaga ggaaccaatg taacgttaca ctgaagaaaa
 4800

cacataataa acggaagaa acggtgtaaa agtgtgaaaa taatTTTTga atatcatttc
4860
ccttggttta attccaaacg aaacgtgtat ttttttagag aatgggaatt cttattggat
4920
5 gtctagattg tttgtttact ccagactgtg cacaaaaacg tttggatgga tgatcagaag
4980
atatTTTTtag gcttagctct aaatataaga aatgatgctt gaaaatccag acagaaattg
5040
10 agtttcaaaa attggaatg tgaggatatta gtcaactaac caaataacaa tgcaaacggg
5100
ttgatacatt tcattttgaa aataatgaaa ctggaattgg atgaccagca cacaaacaca
5160
taaagtaatt atgggaatta gaagcgaaca tagaggaata ctttgccacg aacagaatac
5220
15 aagtgggaac actTTTTtct ccattgtttt agttctgttt ttttgtcaaa ctggttttgt
5280
gctatgtgta aaaaaatatt gccaaagaaa aaagcttggt ttgtggccag tgtccgaaaa
5340
20 aaattttggg gaagcttcgg attaatTTat tttttattc catcggggaa agtggggggg
5400
aaaaaaaaatt taagcagttc ataaaacctt ccaaaaaata tatggacaga gatgattgta
5460
ttttcccgac accaaaatca taattaacta tgagaaaatt gaatgtaacg ttacaattta
5520
25 tttttatttg aagctgaaaa gcgatttatg atttttccga aatgaaaatt ttttttaggt
5580
ttattttttt tgtcgggcaa agaaaaactg aacaaggatt attaaaattt ttggtgtttg
5640
30 ttgtgtctg gagaattcat tcctctctca tcttcacaca atgttttagac atctgacacg
5700
attcaaaata gttcggtttc cgggggttgt gtttagtttt cgtttttcgt tttttttgga
5760
aagaatgttt tagctcattg gttttctttc ttcattcaat agttttgaaa gaatttgccc
5820
35 acttgttatt acaatcatat aaaattaaac tttgatataa aatagagttt gaaagtttcc
5880
cagatccttt ttgatttctt tgtaattttt ttttctccca catatacaca catacaaaacc
5940
gatttttata agaaagagtt ataccctgca gctcgacctc gagggatccg ggccctctag
6000
40 atgcggccgc taggcctcga gggacttttg caccaaaaat aattttattt ccaaaataaa
6060
atttaaataa ataaaaataa ctcataattt aataaaaaatt tcaaaatctt ctagtgtcct
6120
45 ttcatatgca gtacattagc catcagtcac ttaaacagca tctgctgggt gaagaatgct
6180
tgaagcaatt gtccagtccc agaggcacag gctaggagat cttcagtttc ggaggttaacc
6240
tgtaagtctg ttaatgaagt aaaagttcct taggatttcc actctgacta tgggtccaggc
6300
50 acagtgactg tactccttgg ccttcaggta atgcagaatc ctcccataat atcttttcag
6360
gtgcagactg ctcatgagtt ttccccgtgt gaaatcttct ttctccagtt tttcttccag
6420
55 gactgtcttc agatggttta tctgatgata gacattagcc aggaggttct caacaatagt
6480
ctcattccag ccagtgctag atgaatcttg tctgaaaata gcaaagatgt tctggagcat
6540
ctcatagatg gtcaatgagg cgtcctcctt ctggaactgc tgcagctgct taatctcctc
6600
60 agggatgtca aagttcatcc tgtccttgag gcagtattca agcctcccat tcaattgcca
6660

caggagcttc tgacactgaa aattgctgct tctttgtagg aatccaagca agttgtagct
6720
catggaaaga gctgtagtgg agaagcacia caggagagca atttggagga gacacttggt
6780
5 ggtcatgttc ctcgaggcct ttttggccag ctggcgctg ctgcgcgacg gcgagctgct
6840
caccacccag gatccgtccc ctttttcctt tgcgatatc atgtaattag ttatgtcacg
6900
cttacattca cgccctcccc ccacatccgc tctaaccgaa aaggaaggag ttagacaacc
10 6960
tgaagtctag gtccctatct atttttttat agttatgtta gtattaagaa cgttatttat
7020
atttcaaatt tttctttttt ttctgtacag acgcgtgtac gcatgtaaca ttatactgaa
7080
15 aaccttgctt gagaaggctt tgggacgctc gaaggcttta atttgca
7127

United States Patent & Trademark Office
Office of Initial Patent Examination -- Scanning Division



Application deficiencies found during scanning:

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not
present
for scanning. (Document title)

* Scanned copy is best available. page 71 - page 64 as part
of specification are
Sequence listing